

Distress in cancer Patients

by D Denys

Submission date: 30-May-2018 10:39PM (UTC+0500)

Submission ID: 970382743

File name: Distress_in_Cancer_Patients...rtf (112.15K)

Word count: 2339

Character count: 13830

Distress in Cancer Patients

Name

Institutional Affiliations

Distress in Cancer Patients

Cancer is often perceived to be **untreatable**, an unbearable painful disease with no cure. Having a full awareness of the long-term psychological and intermediate repercussions of inherited trials for patients with cancer motivates research, unfortunately, an open agreement of the psychological, social effects and check-up regularity for cancer-affected *BRCA 1 and BRCA 2* alteration transferors is not available. The aim of this paper is to discuss various methods of treating cancer and taking care of patients with cancer.

Systematic review of prolonged and intermediate studies was carried out testing the psychological effects such as depression, anxiety and psychological distress among the affected in comparison to the unaffected alteration carriers. The evaluation involved the repeat of several studies. The majority of long-term and intermediate research dwelling on depression and other symptoms of distress and anxiety among the affected was employed. **This was basically done on the discovery of the testing results aided by the carrier factors.** The investigation established a number of patient outcomes, designs and methods. Evidence showing that victims of cancer and mutations encountered undesirable effect related to mental uprightness in reference to rising in distress symptoms, depression, and anxiety in the initial months after the disclosure test (Ringwald, Johanna and et al., 2016).

According to Mu'taz, Asmar and Al-Rjoub (2016), cancer patients are exposed to health problems and emerges as symptoms and signs which are mental and bodily in nature. Moreover, increased ranks of hopelessness and poor health conditions were shared among the patients. These problems are related to the impact on the cancer **patients'** emotional, spiritual well-being, psychological, social, and physical that eventually deteriorate QoL (Quality of Life) of patients.

Furthermore, depression and fatigue also affect the outcomes of patients treated. Fatigue being a multidimensional symptom, affects the cancer patients' life significantly. In addition, ¹⁵ 66% of patients with the breast related cancer suffered from exhaustion after testing. Also, this condition went beyond treatment end. Patients with cancer who begins treatment, experiences fatigue and later encounters worse symptoms.

Da Mata, Luciana Regina Ferreira and et al (2016), states that surgery is the principle technique of management and it is used mostly in patients diagnosed with cancer. Patients view surgical processes as being aggressive due to uncertainty. Moreover, fear of physical body change causes stress and eventually conflict situations; women who undergo partial mastectomy surgery experiences stress and differences in self-confidence due to the phobia of losing sexuality and femininity symbol. In addition, these experiences lead to low self-esteem which is a determining factor that ensures the psychological and physical well-being, which eventually has a positive or negative influence on the mode of dealing with the disease and the corresponding treatment. Nurses can intervene on realizing the existence of sorrow and reduced self-confidence. This can also be done in the assistance of other interdisciplinary members within the units, mostly in the evaluation period, with a goal of reducing the impact brought by stress. However, this article hypothesized those cancer patients, in a time of 48hours of mediate postoperative displays these changes in self-esteem and distress.

In relation to Mosleh, Sultan and et al. (2018), low self-confidence and nervousness are common symptoms possessed by cancer patients in Jordan. These results were found out when an expressive design method was employed to conduct research. 226 cancer patients completed the social difficulties inventory, EORTC-QoL-C30, comfort scale and Sickbay unease and

misery scale (HAD). Mosleh, Sultan, et al. (2018), found out that Hopelessness was the primary responsive problem with a greater occurrences compared to nervousness (67.6% vs. 43%). Patients with cancer experiencing the two symptoms encountered inferior QoL scores and high marks for protests in relation to symptoms. The cancer patients remained possible to undergo public difficulties in their daily life. In addition, older people with cancer displayed high depression, social difficulty and poor QoL. On the other hand, anxiety was high in female patients in advanced cancer stage, poor QoL and hospital readmission.

Cancer patients ought to accept and move on with life after diagnosed with cancer and navigate a fragmented healthcare system. Patient navigation is of great help to family members and the cancer patients since it provides emotional and tangible assistance. Patients' navigators guide patients through the unknown slope of modern cancer care, attending to the patient's questions and being available when needed by the patients. Furthermore, navigation programs are characterized with providing patient-centered guidance and support with an aim of motivating the patients to overcome challenges like on-time diagnosed resolution, engagement of patient in health-care activities, the satisfaction of patient and cost-effective care (Cobran, Ewan and et al. 2017).

According to Gundelach, Amy, and Barb Henry (2016), patients with cancer are exposed to psychological distress. Psychological distress is a negative emotion which interferes with daily activities and affects individual's level of functioning. In addition, psychological distress arises from negative perceptions of the self and circumstances and is portrayed by anxiety, sadness, symptoms of mental illness, and distraction. Moreover, psychological distress has an impact on family, spirituality and finances. Krebber, Marie and et al. (2016) found out that it is a challenge

to recognize and support patients of cancer suffering from psychological agony specifically, those affecting the head and the neck (HNC). However, screening is efficient at the subdivision of Otolaryngology, which focuses on head and neck surgery, followed by triage to care since 2016 by a trace shade computer scheme. The computer scheme is associated with the information of patients in the hospital. The process of records is faster because the nurse specialized in the Head and neck division looks at the patients' results simply presented in illustrations and later discusses it the patient.

Ng, Chong Guan and et al. (2015), highlighted that breast cancer was a common disease among females in Malaysia. In the past, breast cancer patients' quality of life improved due to early detection and early treatment. Eminence of Life is the perception of the well-being of a patient. Quality of life details revolve around cognitive, mental, social and physical functioning. In addition, nervousness and despair are the main collective psychiatric challenges suffered by victims of breast cancer. Patients with breast cancer encounter depression across the stages of their illness. Research done on anxiety and depression among these patients in Malaysia established that anxiety levels were at 31.7% and depression at 22.0%. Factors such as being the single, financial burden, and young age contributed significantly to anxiety among breast cancer patients in Malaysia. According to Nottelmann, Lise and et al. (2017), patients with cancer face unfulfilled palliative care; the physical, social, emotional and cognitive challenges. Comforting care is applied immediately when the illness is detected together with therapies. However, soothing care is associated with losing hope among the caregivers and cancer patients. Ultimately, these people are not referred to particular palliative maintenance early till the point where the illness is trailed. In accordance to The World Health Organization (2002), intensive

care should be provided to families as well as all patients with this threatening illness. This should mainly be a cohesive treatment in the early phase of cancer. Furthermore, rehabilitation is important since it is involved in refining and upholding mental, intellectual and social enactment levels, physical and prevents retard of roles which relates to daily activities.

The cognitive element of the emotional experience of anxiety is a worry. Worry is the main symptom of generalized anxiety disorder (GAD). It focuses on the future problems and how to tackle the problem. In addition, the worry is associated with various adverse effects. This is related to depression and anxiety than contemplation. Also, it mediates the relationship between personality factors like the autonomic nervous system, increased cortisol output, elevated interleukin-6 responses prior to the influenza vaccination, and susceptible disturbances following the introduction to a dreaded stimulus. (Wu, Salene and et al. 2013)

In reference to Whitaker and et al. (2016), patients' emotional responses to symptoms influence behavioral responses. The symptom appraisal process is the main step to achieving earlier presentation, and the earlier diagnosis of cancer which leads to better outcomes. Elefteria (2017), states that depression is a serious and common mental disease currently. Depressive disorders are common among single people. Also the depression prevalence rates from 2% for a month to 6% from the rest of patients' life. Apart from depression, patients with cancer undergo pain as death approaches. In addition, depression and pain have an effect on thoughts, moods, and is accompanied by the disturbance of the person's eating habits and sleep, and bodily discomfort. According to Bseiso, Reema Awni, & Abdelaziz Mousa Thabet 2017, there has been an increasing number of cancer patients in Palestine. Breast cancer was ranked top and was highest in women and focused on the age group of twenty to fifty-nine. Large intestine cancer

was the second according to (MOH, 2014) report. In the report of MOH, cancer is another major cause of death in Palestine. It was impossible for these patients to receive treatment in Israel or Cairo simply due to the closure of the borders. War and siege activate these stressors and raises death mortality due to limited treatments, transfer procedures and equipment.

Jackson and et.al (2017) indicated that levels of psychological distress are high because of reduced health-related quality of life (QOL). Patients that are psychologically distressed seek medical services, are less adherent to their treatment regimens, experience difficulty in making decisions, and less satisfied with their medical care. In the end, the patients experience poor QoL, unnecessary emotional suffering and emotional vulnerability. There is a need for psycho-oncological education to cancer patients regardless of the illness trajectory. According to Butow, Phyllis and et al. (2015), medical trails offer evidenced recommendations to aid best preparation and reliable care for certain patient apprehension in homogeneous patient group. Clinical pathways have effectively increased uncovering of emotional indispositions and rates of recommendations for treatment. Finally, despite the fact that there has been a tangible advancement in the levels fight against this enemy, there are still a lot of mileages to be achieved concerning the costs and availability of respective treatments. The overall costs outstanding are somehow expensive due to the inadequate nature of the prevailing infrastructural facilities.

In conclusion, innovations in cancer care and approaches are necessary to increase access and improve affordability. The palliative program should be implemented in order to allow patients with progressive get adequate relief from their spiritual, emotional and physical suffering. Again, this should adequately be done to benefit those who have little access to the treatment mode. It is much evidenced that newly diagnosed patients with cancer will only have a

decreased anxiety and distress specifically, when mentored with other cancer survivors or through participation in other partnering programs. Moreover, participation in this peering programs and in adherence to treatment plans under mentoring programs for a period of about six weeks is likely to curb the discussed impacts.

References

Bseiso, R. A., & Thabet, A. M. (2017). The Relationship between Siege Stressors, Anxiety, and Depression among Patients with Cancer in Gaza Strip. *Health Science Journal*, 11(2).

Butow, P., Price, M. A., Shaw, J. M., Turner, J., Clayton, J. M., Grimison, P., ... & Kirsten, L. (2015). Clinical pathway for the screening, assessment and management of anxiety and depression in adult cancer patients: Australian guidelines. *Psycho-Oncology*, 24(9), 987-1001.

Cobran, E. K., Merino, Y., Roach, B., Bigelow, S. M., & Godley, P. A. (2017). The Independent Specialty Medical Advocate Model of Patient Navigation and Intermediate Health Outcomes in Newly Diagnosed Cancer Patients. *Journal of Oncology Navigation & Survivorship*, 8(10).

da Mata, L. R. F., da Silva, M. R., Antunes, A. C. C., Faria, B. S., Chávez, G. M., & Oliveira, P. P. (2016). Self-esteem and distress in patients undergoing cancer surgery: a correlational study. *Online Brazilian Journal of Nursing*, 15(4), 664-74.

Gundelach, A., & Henry, B. (2016). Cancer-Related Psychological Distress: A Concept Analysis. *Clinical journal of oncology nursing*, 20(6).

Hendricks-Jackson, L., Smith, K., Kressel, B., Bishop, C., Thornton, K., Posmontier, B., & Paller, C. J. (2017). Psycho-Oncological Education to Reduce Psychological Distress Levels in Patients with Solid Tumor Cancers: A Quality Improvement Project. *Journal of Oncology Navigation & Survivorship*, 8(10).

- Krebber, A. ⁷ M. H., Jansen, F., Cuijpers, P., Leemans, C. R., & Verdonck-de Leeuw, I. M. (2016). Screening for psychological distress in follow-up care to identify head and neck cancer patients with untreated distress. *Supportive Care in Cancer*, 24(6), 2541-2548.
- ⁹ Mosleh, S. M., Alja'afreh, M., Alnajjar, M. K., & Subih, M. (2018). The prevalence and predictors of emotional distress and social difficulties among surviving cancer patients in Jordan. *European Journal of Oncology Nursing*, 33, 35-40.
- ¹¹ Mu'taz, M. D., Asmar, I. T., & Al-Rjoub, B. (2016). An Original Research about: The Associations of Depression and Fatigue with Quality of Life among Palestinian Patients with Cancer. *Health Science Journal*, 10(6).
- ³ Ng, C. G., Mohamed, S., See, M. H., Harun, F., Dahlui, M., Sulaiman, A. H., ... & Taib, N. A. (2015). Anxiety, depression, perceived social support and quality of life in Malaysian breast cancer patients: a 1-year prospective study. *Health and quality of life outcomes*, 13(1), 205.
- ⁶ Nottelmann, L., Groenvold, M., Vejlgard, T. B., Petersen, M. A., & Jensen, L. H. (2017). A parallel-group randomized clinical trial of individually tailored, multidisciplinary, palliative rehabilitation for patients with newly diagnosed advanced cancer: the Pal-Rehab study protocol. *BMC cancer*, 17(1), 560.
- ¹³ Papamichail Eleftheria MSc, R. N. (2017). Depression, Pain and Mental State in Cancer Patients. *International Journal of Caring Sciences*, 10(2), 955.

⁸ Ringwald, J., Wochnowski, C., Bosse, K., Giel, K. E., Schäffeler, N., Zipfel, S., & Teufel, M. (2016). Psychological distress, anxiety, and depression of cancer-affected BRCA1/2 mutation carriers: A systematic review. *Journal of genetic counselling*, 25(5), 880-891.

¹⁰ Wu, S. M., Schuler, T. A., Edwards, M. C., Yang, H. C., & Brothers, B. M. (2013). Factor analytic and item response theory evaluation of the Penn State Worry Questionnaire in women with cancer. *Quality of Life* ¹² Whitaker, K. L., Cromme, S., Winstanley, K., Renzi, C., & Wardle, J. (2016). Emotional responses to the experience of cancer 'alarm' symptoms. *Psycho-Oncology*, 25(5), 567-573. *Research*, 22(6), 1441-1449.

Distress in cancer Patients

ORIGINALITY REPORT

24%

SIMILARITY INDEX

14%

INTERNET SOURCES

10%

PUBLICATIONS

23%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Grand Canyon University

Student Paper

4%

2

Submitted to 97136

Student Paper

2%

3

Submitted to University of Strathclyde

Student Paper

2%

4

Submitted to Trinity College Dublin

Student Paper

2%

5

Submitted to University of New England

Student Paper

2%

6

Submitted to University of Louisiana, Lafayette

Student Paper

2%

7

Submitted to University of Hertfordshire

Student Paper

2%

8

jultika.oulu.fi

Internet Source

2%

9

Submitted to La Trobe University

Student Paper

2%

10	www.vpgcentral.com Internet Source	1%
11	Submitted to Roane State Community College Student Paper	1%
12	Submitted to Australian Catholic University Student Paper	1%
13	Submitted to DeVry, Inc. Student Paper	1%
14	Submitted to The University of Notre Dame Student Paper	<1%
15	www.hsj.gr Internet Source	<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

Distress in cancer Patients

PAGE 1

PAGE 2



Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.



Possessive This word may be a plural noun and may not need an apostrophe.

PAGE 3



Possessive You may need to use an apostrophe to show possession.

PAGE 4



Wrong Article You may have used the wrong article or pronoun. Proofread the sentence to make sure that the article or pronoun agrees with the word it describes.



Article Error You may need to remove this article.



Article Error You may need to use an article before this word.



Article Error You may need to remove this article.



Prep. You may be using the wrong preposition.



Article Error You may need to use an article before this word.



Article Error You may need to use an article before this word.



Article Error You may need to use an article before this word.



Article Error You may need to use an article before this word.



Pronoun This pronoun may be incorrect.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.



Article Error You may need to use an article before this word. Consider using the article **the**



Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.



Frag. This sentence may be a fragment or may have incorrect punctuation. Proofread the sent to be sure that it has correct punctuation and that it has an independent clause with a complete subject and predicate.

PAGE 5



Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.

PAGE 6



Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.



P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.

PAGE 7



Article Error You may need to use an article before this word. Consider using the article **the**



Article Error You may need to use an article before this word.

PAGE 8

PAGE 9

PAGE 10

PAGE 11
