

Exploring Theories of Victimization Using a Mathematical Model of Burglary

Journal of Research in Crime and

Delinquency

48(1) 83-109

© The Author(s) 2011

Reprints and permission:

sagepub.com/journalsPermissions.nav

DOI: 10.1177/0022427810384139

<http://jrcd.sagepub.com>



Ashley B. Pitcher¹ and Shane D. Johnson²

Abstract

Research concerned with burglary indicates that it is clustered not only at places but also in time. Some homes are victimized repeatedly, and the risk to neighbors of victimized homes is temporarily elevated. The latter type of burglary is referred to as a near repeat. Two theories have been proposed to explain observed patterns. The boost hypothesis states that risk is elevated following an event reflecting offender foraging activity. The flag hypothesis, on the other hand, suggests that time-stable variation in risk provides an explanation where data for populations with different risks are analyzed in the aggregate. To examine this, the authors specify a series of discrete mathematical models of urban residential burglary and examine their outcomes using stochastic agent-based simulations. Results suggest that variation in risk alone cannot explain patterns of exact and near repeats, but

¹ Centre d'Analyse et de Mathématiques Sociales, Ecole des Hautes Etudes en Sciences Sociales, Paris, France

² UCL Jill Dando Institute of Crime Science, University College London, London, United Kingdom

Corresponding Author:

Shane D. Johnson, UCL Jill Dando Institute of Crime Science, University College London, Second Floor Brook House, 2-16 Torrington Place, London, WC1E 7HN, United Kingdom

Email: shane.johnson@ucl.ac.uk

that models which also include a boost component show good qualitative agreement with published findings.

Keywords

burglary, mathematical model, repeat victimization, boost hypothesis, risk heterogeneity, agent-based simulation

Introduction

Criminology is a multidisciplinary subject but the application of mathematics has been fairly limited in the study of crime patterns (for recent exceptions, see Short et al. 2008; Short et al. 2010; and, for a special journal issue devoted to this, see the *European Journal of Applied Mathematics*, Volume 21, Issues 4-5). The advantages of using a mathematical approach are numerous and range from the conceptual clarity that arises from using formal expressions to describe systems, to the ability to model nonlinear complex systems, to the possibility of developing control strategies if systems are adequately specified. In this article, we use mathematical models to examine theories concerned with space-time patterns of victimization at the level of place and consider how concepts commonly used in mathematics may be helpful in formalizing and exploring theories of this kind.

Whatever the method, the unit of analysis selected will directly affect the reliability of inferences drawn. For example, where research concerns theories of crime concentration, the analysis of data aggregated to large geographical units will often be inappropriate and invite errors of inference. A common example is the ecological fallacy (Robinson 1950) of assuming that patterns observed in the aggregate across an area will apply to all places within it. Rarely will this be the case. For example, Bowers, Johnson, and Pease (2005) demonstrate that for domestic burglary irrespective of what type of area they live in, the type of house (e.g., flats, row homes, and so on) a resident lives in significantly affects their risk of victimization (see also, Budd 1999). Likewise, a series of studies demonstrate that regardless of area-level risks, how connected a block face is to the surrounding street network significantly influences the risk of burglary to homes located on it (e.g., Beavon, Brantingham, and Brantingham 1994; Bevis and Nutter 1977; Hillier 2004; Johnson and Bowers 2010). Stated more explicitly, the importance of place in the study of crime—where places may include meaningful units of analysis such as block faces or individual homes—has been acknowledged for some time (e.g., Eck and Weisburd 1995).

However, location is not the only dimension of importance in the study of crime patterns. Crimes occur at a point in time as well as a place, and to neglect one aspect may lead to an inadequate understanding of the contribution of the other. For example, where the dimension of time is ignored the stability over time of factors that affect crime risk at places may be overestimated. In line with this, while it has been established for some time that crime clusters in space at a range of spatial scales from areas to micro-level places (e.g., Pease 1998; Ratcliffe 2004; Shaw and McKay 1969; Sherman, Gartin, and Buerger 1989; Weisburd et al. 2004), research shows that it clusters in *both* space *and* time (e.g., Grubestic and Mack 2008; Johnson et al. 2007; Townsley, Homel, and Chaseling 2003). For example, at the finest spatial scale, numerous studies (for a review, see Farrell 2005) have shown that some homes are repeatedly victimized more often than would be expected on a chance basis, assuming the risk of crime were uniform. Moreover, when repeat burglary victimization occurs, it is more likely to do so swiftly than after some time has elapsed. In fact, the time course of repeat victimization fits an exponential decay function rather well (Johnson, Bowers, and Hirschfield 1997; Polvi et al. 1991; Townsley, Homel, and Chaseling 2000). More recent work (e.g., Grubestic and Mack 2008; Johnson et al. 2007; Johnson, Summers, and Pease 2009; Short et al. 2009; Townsley et al. 2003) suggests that this phenomenon extends to nearby homes such that when one house is victimized, those nearby also appear to experience a temporary elevation in risk. When this occurs, it has been referred to as a *near repeat* (Morgan 2001; Townsley et al. 2003).

With respect to repeat victimization of the same home, places would be defined as individual households (Eck and Weisburd 1995). However, this has the potential to imply that individual homes might be considered independent units for which the risk of victimization is unaffected by features of their neighbors. That neighbors of victimized homes experience a temporary elevation in risk following an offense implies a dependency and that places may be better conceptualized as the slightly larger spatial units such as block faces, or small clusters of homes that immediately surround each individual housing unit. Alternatively, perhaps specifying places as individual homes *is* most appropriate but what is required is a better understanding of how particular places influence the risk of crime to others nearby. Thus, it would seem that further research is required to better understand how risk varies at the level of place and for the testing or refinement of theories that might explain observed patterns.

Two theories have been proposed to explain patterns of repeat victimization and by association, near repeats. According to the *flag hypothesis*, observed patterns can be explained in terms of (relatively) time-stable

variation in risk heterogeneity across units (Nelson 1980; Sparks 1981). This variation in risk may be influenced by a variety of factors including those already discussed, such as accessibility and target attractiveness. These may vary at both the individual household, block face and neighborhood level, suggesting that the risk of victimization at places may be influenced by the time-stable characteristics of the spaces within which they are located as well as features of the places themselves.

An alternative focus suggests that the risk of victimization at places varies (to some extent) over time as a function of *current* patterns of victimization. Simply put, following one victimization the risk of crime is said to be temporarily *boosted* (Pease 1998) to victimized homes and those nearby (Johnson and Bowers 2004; Johnson et al. 2007; Johnson et al. 2009; Townsley et al. 2003), most likely reflecting dynamic foraging strategies on the part of offenders (Johnson and Bowers 2004; Johnson et al. 2009). Thus, risk is considered a function of changes in offender awareness (e.g., see Brantingham and Brantingham 1993, 2008) and perceptions of crime opportunities at particular places within spaces.

As discussed by Johnson (2008), taken on face value, the time course of repeat victimization would appear to support the boost account of repeat victimization; if the risk of victimization was time stable why would an exponential decay in risk be observed? However, this perception may be illusory and instead reflect a type of statistical artifact that can occur when data are aggregated for groups of places that experience very different risks (for a more general discussion of heterogeneity's ruses, see Vaupel and Yashin 1985). The potential problem is that the researcher may erroneously assume that aggregate patterns reflect those for individual places. To illustrate, consider an area in which there are three classes of home; those with stable-high, -medium and -low risks of victimization. Even on a chance basis, some homes from each group would experience repeated victimizations. If these occurred purely by chance—insofar as they were unrelated in terms of who committed the offenses—and analyses were performed independently for each class of home, the time course of repeat victimization would be uniform over time for each group. However, if the data for the three groups were combined, a curve is likely to be generated. To elaborate, the high-risk homes would be victimized the most and would be the most likely to be revictimized swiftly. Considering the low-risk group, they would be victimized least often but might still be revictimized. When they are, the time to revictimization would typically be longer than that for the other two groups. The medium-risk group would experience repeat victimization at a rate somewhere between the other two groups. Mixing the patterns for the

three groups would generate a curve for which the coefficient would be a function of the differences in risk for the three populations; the greater the differences, the more accentuated the curve. Thus, risk heterogeneity can plausibly explain the observed time course of repeat victimization. The same explanation can be extended to patterns of space–time clustering more generally (near repeats) if there is sufficient variation in risk across—but homogeneity within—neighborhoods or (for example) street segments.

To examine these theories, a number of research methods can be and have been employed. For example, to examine repeat victimization, multivariate statistical methods have been used to estimate the extent to which patterns of concentration observed in cross-sectional data can be explained by measured heterogeneity across homes (Osborn and Tseloni 1998). Such analysis suggests that heterogeneity explains some but not all of the variation in concentration observed. Analyses of crimes detected by the police suggest that crimes committed at the same location or nearby are likely to be the work of the same offender/offenders (e.g., Bernasco 2008; Johnson et al. 2009; Kleemans 2001), and interviews with offenders provide further support for both the boost (e.g., Ashton et al. 1998; Ericsson 1995) and flag (e.g., Rengert and Wasilchick 2000) explanations.

However, criticisms of these approaches suggest that the use of other complementary methods would enhance criminological understanding. For example, as Eck and Liu (2008a) discuss, detection rates are so low that while analyses of such data are important, it is possible that the data are biased, reflecting the behavior of a limited sample of offenders, who are (for example) perhaps those easiest to apprehend. A similar critique can be applied to much of the ethnographic research, as offenders are usually identified through contact with the criminal justice system and thus only represent those known to authorities.

In contrast, where response rates are high, data obtained from large-scale surveys may be relatively unbiased and provide a good estimate of actual patterns. Unfortunately, data collected using sample surveys will rarely (if ever) provide detailed data that allow the analysis of micro-level spatial and temporal patterns of crime. For crimes recorded by the police, there will generally be issues with underreporting (e.g., Xie et al. 2006), but for some types of crime (such as burglary with loss) this may be less of a problem than for other types of crime. However, for both types of data (survey and police recorded crime data), the types of analysis conducted essentially rely on correlation as an approach to hypothesis testing. That is, data for a series of independent variables—intended to represent theoretical constructs of interest—are collected and a statistical model used to estimate the degree

to which they are associated with the dependent variable of interest. While fairly sophisticated models are available, that can be used to estimate and control for the influence of measured and unmeasured variables, correlation does not imply causation and nor does it help to specify the precise mechanism/mechanisms through which outcomes are generated (see Eck and Liu 2008a).

In contrast to inductive approaches, mathematical and simulation models are used to formally specify a theory and to then evaluate whether the model can generate patterns which resemble “statistical signatures”¹ or even precise outcomes (Gilbert 2008). This type of approach has a number of merits to recommend it. We will not discuss all of these here, as they are dealt with elsewhere (Eck and Liu 2008b), but some points are as follows. First, such models require that the precise mechanisms through which a pattern is thought to emerge are explicitly stated, along with assumptions made. This formal specification requires a level of precision that can be absent in theories expressed in spoken languages (Eck and Liu 2008a), allowing the internal logic of the theory to be tested. This is, in and of itself, a useful exercise. Second, such models allow theory falsification; if a model cannot generate a pattern then the theory can be considered insufficient (Eck and Liu 2008a). Third, such models can generate data for analysis, but unlike other research methods, models can be executed numerous times allowing outcomes to be examined for consistency or allowing the effects of particular manipulations to be observed. Thus, mathematical models may provide a useful additional method of theory testing, particularly where representative data are difficult to obtain.

For such reasons, using a simple stochastic simulation model, Johnson (2008) examined the plausibility that the time course of repeat victimization might be explained by a ruse of heterogeneity (see also Short et al. 2009) but concluded that this alone could not explain observed patterns. This is important but no research of which we are aware has examined this for more general patterns of space–time clustering (i.e., near repeats) and hence this is the focus of the current article. Moreover, in the previous studies, because of the ways the models were specified, the risk to each home was essentially independent of every other. In reality, it may be the case that the risk of victimization to any home is a function of the target attractiveness of that place but also of those located nearby. For example, while a home of low attractiveness might be at little risk of victimization when considered independently, it may be at a greater risk if it is surrounded by homes that represent good targets for burglary, which might attract the attention of offenders already in the locale.

In this article, our goal is to examine the potential contributions of the two theoretical mechanisms in the generation of space–time patterns of burglary victimization using a simple mathematical model which takes account of the fact that the risk to one home may be partly a function of the target attractiveness of those located nearby. Our aim is not to reproduce observed patterns precisely but rather to test the more general middle-range theories (see Gilbert 2008) described. As such, in what follows we do not compare our results to those observed for a particular area but to the statistical signatures summarized above.

In what follows, we consider a home's attractiveness to burglary as the statistical rate of burglary when a burglar is present, as described in Short et al. (2008). We will describe a modified version of their mathematical model that will govern the dynamics of the attractiveness of places and burglar movement. Stochastic agent-based simulations will then be used to generate burglary data that can be checked for qualitative agreement with observed patterns reported in the existing research literature.

The Mathematical Model

According to the boost hypothesis, a burglary event causes the risk of further victimization to be elevated at the burgled house and those nearby. This elevation in risk decays in time and as distance from the burgled house increases.

We consider a uniform square lattice with one home located at each lattice site s and characterize each home by its attractiveness to burglary, measured by the sum of two variables: $\tilde{A}_s$ and B_s^t , where $\tilde{A}_s$ is a static component and B_s^t is a dynamic component for the time period $(t, t + \delta t)$, where time is measured in days. The static component $\tilde{A}_s$ represents the risk from factors associated with places that do not tend to change over a short time-scale, and the dynamic component B_s^t captures the boost to burglary risk caused by previous burglaries. The total attractiveness of a home at site s for the time period $(t, t + \delta t)$ is then given by:

$$A_s^t = \tilde{A}_s + B_s^t, \quad (1)$$

The total attractiveness A_s^t can be thought of as the rate of burglary when a burglar is present. Burglary is assumed to follow a Poisson process with rate parameter A_s^t , so that the mean number of burglaries during the time period $(t, t + \delta t)$ is $A_s^t \delta t$. When δt is small, we can assume that only one burglary event can occur per time step. This means that the probability a burglar at

house s commits a burglary during the time period $(t, t + \delta t)$ can be simplified to $1 - \exp(-\tilde{A}_s \delta t)$.

We update the dynamic attractiveness after each time step, according to a stochastic difference equation. To formulate this equation, we assume that a proportion of the dynamic attractiveness $\left(\frac{\eta}{4}\right)$ at a site s moves to each of its four neighbors after each time step and that the proportion $(1 - \eta)$ of the dynamic attractiveness remains at site s . The parameter η is used to model the rate of spatial diffusion per unit time; the larger the value of η , the faster the risk diffuses. According to the boost hypothesis, dynamic attractiveness should also fade over time, and hence a multiplicative decay term $(1 - \omega \delta t)$ is included, where δt is the time-step size in days and $1/\omega$ is the mean lifetime of dynamic attractiveness. Every time there is a burglary at a site s , dynamic attractiveness increases by $\theta \left(1 - \frac{B_s^t}{k}\right)$, where θ describes the magnitude of the boost effect per burglary and k is a measure of carrying capacity for dynamic attractiveness. The inclusion of a parameter to model carrying capacity ensures that the attractiveness of each home does not exceed an unrealistic level. Expressed another way, if the risk to a home continues to increase, we assume that there will be a tipping point after which the risk of victimization will decrease (until it is back to the tipping point). Carrying capacity may be thought of as reflecting the effects of a police response to an observed elevation in crime concentration in a neighborhood (or at a specific place), defensive behavior on the part of the residents, or offenders' anticipation of one or both of the two. To model the number of burglaries that occur at a site s , we let X_s^t be a random variable for the number of burglaries that occur at site s during the time period $(t, t + \delta t)$. If there is no burglar at a site s during the time period $(t, t + \delta t)$ then X_s^t takes a value of zero. Otherwise, it is Poisson distributed with mean $A_s^t \delta t$, that is, $X_s^t \sim \text{Poisson}(A_s^t \delta t)$. This leads to the stochastic difference equation:

$$B_s^{t+\delta t} = \left((1 - \eta)B_s^t + \frac{\eta}{4} \sum_{s' \in os} B_{s'}^t \right) (1 - \omega \delta t) + \theta X_s^t \left(1 - \frac{B_s^t}{k} \right) \quad (2)$$

Note that $\omega \delta t$ must be between zero and one to be sensible. The notation $s' \in os$ refers to all the sites adjacent to site s , which according to the assumptions in this model, are the four nearest neighbors on the square lattice (see Figure 1). Equation (2) is similar to equation (2.5) in Short et al. (2008)² but adds the carrying capacity term and the stochastic random variable X_s^t . The carrying




Figure 1. Four nearest neighbors on the square lattice (Von Neumann neighborhood).

capacity term k is important for theoretical reasons, but it also limits the sensitivity of the model to changes in parameter values. For example, an increase in θ causes the number of burglaries to increase, but the carrying capacity term ensures that the total number of burglaries will be within a reasonable bound and hence that the volume of crime does not go off to infinity.

In order to determine where burglaries occur, it is necessary to define the rules that govern the location and movement of burglars on the lattice. These are summarized in Figure 2 and articulated in a little more detail here. We assume that burglars are generated at particular locations by way of a Poisson process with rate parameter that is the sum of a spatially uniform component (Γ) and a component dependent on target attractiveness (βA_s^t). In other words, we generate burglars on the lattice according to a Poisson process with mean $(\Gamma + \beta A_s^t)\delta t$, where Γ and β are positive constants.

We assume that burglars tend to move toward houses with higher attractiveness. To model this, for each time step and each burglar located at house s , we calculate the following movement probabilities:

$$\rho_{right} = A_{right}^t / T_s^t \quad (3a)$$

$$\rho_{up} = A_{up}^t / T_s^t \quad (3b)$$




Figure 2. Overview of the algorithm used in the simulation.

$$\rho_{left} = A_{left}^t / T_s^t \quad (3c)$$

$$\rho_{down} = A_{down}^t / T_s^t \quad (3d)$$

where the subscripts indicate the nearest neighbor to s on the square lattice, and T_s^t is the sum of the attractiveness of all the nearest neighbors to house s for time period $(t, t + \delta t)$, that is,

$$T_s^t = \sum_{s'os} A_{s'}^t. \quad (4)$$

The four movement probabilities sum to one, that is,

$$\rho_{right} + \rho_{up} + \rho_{left} + \rho_{down} = 1. \quad (5)$$

We then sample a random number, call it r , from the uniform distribution on the interval $[0,1]$ so that the rules of burglar movement are:

if $r \leq \rho_{right}$	$\Rightarrow$ move right	(6a)
--------------------------	--------------------------	------

else if $r \leq \rho_{right} + \rho_{up}$	$\Rightarrow$ move up	(6b)
---	-----------------------	------

else if $r \leq \rho_{right} + \rho_{up} + \rho_{left}$	$\Rightarrow$ move left	(6c)
---	-------------------------	------

Else	$\Rightarrow$ move down	(6d)
------	-------------------------	------

Burglars can only move once per time step, so the size of the time-step δt tells us how fast burglars move from house to house. We have used $\delta t = 1/2000$, meaning that burglars move every 43.2 sec. This gives each burglar the chance to assess the potential target.³ As a result of the use of a biased random walk, the risk to each home is influenced by its own attractiveness and that of its neighbors.

Burglars eventually get tired and leave the lattice after f days (the “fatigue time”). For simplicity in the current study, we have used $f = 1/24$ (or 1 hr) for all agents. Other possibilities, such as varying this parameter across agents, are acknowledged but not examined here.

It is important to note that this model was designed to examine patterns of crime at the level of place. It is not unit free and cannot be realistically applied to very different spatial scales. For example, each site s cannot represent a unit such as an entire county because burglars cannot evaluate the relative attractiveness of neighboring counties and cannot travel quickly between them; but they can do so with neighboring houses. As a general point, very rarely will the same model be valid for all spatial scales. In fact, an entire branch of mathematical modeling—multiscale modeling—is devoted to understanding large complex systems by linking together different models on different spatial scales. A recent example of this is Shipley and Chapman (2010), where multiscale modeling is used to link different models for fluid and drug transport in vascular tumors. The model on the capillary length scale is different from the model on the tumor length scale, and special mathematical techniques are used to link the two models together. We do not apply such techniques here, but their consideration may be useful in future research concerned with how factors that operate at different spatial scales influence crime at places.

Counting Repeats and Near Repeats

For n burglaries (where $n > 1$) occurring at a house s , each of the possible $n(n - 1)/2$ burglary pairs are counted as repeats. For example, consider three burglaries that all occur at house s . The first burglary occurs on day 1, the second on day 3, and the third on day 6. In this case, the number of repeat burglaries at house s is three and the times between repeats are 2 days (between the first and second burglary), 3 days (between the second and third burglary) and 5 days (between the first and third burglary).

The alert reader may ask why we do not simply count the $(n - 1)$ intervals between sequential events? The reason for this becomes clearer when

considering near repeats. In this case, when a burglary occurs close in space and time to more than one other, we would have to generate a clear set of rules for determining which of the previous burglaries should be counted for the purpose of determining the space and time between near repeats. Should one favor closer distance in determining near repeats or a shorter time between events? It would be difficult to give an objective answer to this question, so we instead consider every possible burglary pair.

There is also a counting issue with the square lattice. When a burglary occurs, there are more homes $n + 1$ doors away than n doors away. For example, when a burglary occurs, there are four homes one door away but eight homes two doors away (diagonal moves are counted as two doors away in a Von Neumann neighborhood). To account for this, one solution is to divide the number of burglaries that occur n doors away from a previously burgled home by $4n$ for $n \geq 1$. Failing to standardize the results by opportunity in this way would confound interpretation of the results. Accordingly, the vertical axes in all graphs that follow are labeled "Number of burglaries (adjusted)."

However, there is also an issue with using a finite spatial domain in the model since places near the boundary do not have as many close neighbors as those further away from it; a spatial edge effect (Boots and Getis 1988). The influence of the edge effect is a function of two things. First, it will be inversely proportional to the size of the spatial domain. Second, it will be a function of the distance over which effects are measured. For example, if we only considered events that occurred within two homes of each other, the edge effect would be much less pronounced than if we examined those for homes 100 doors apart. To correct for this, we derived a general equation for calculating the mean number of opportunities in an $N \times N$ grid that are n homes apart⁴ which can be used to standardize the results:

$$\text{mean}(n, N) = \frac{4}{N^2} \left(n(N - 2n)^2 + \sum_{k=n+1}^{2n} [k(k - n) + (2k - 1)(N - 2n)] + 2 \sum_{k=n+1}^{2n-1} [k(2n - k)] \right) \quad (7)$$

To illustrate the scale of the edge effect (which is quite small) for a grid of 100×100 homes, for homes that are four doors apart, rather than standardizing the results by dividing the number of observed events by $4n$ (16),

Table 1. Parameter Values for the Mathematical Model

Parameter	Value
δ_t	1/2000
η	1/5000
θ	0.18
κ	0.18
ω	1/42
Γ	1/100
β	5
f	1/24

we divide by 15.36. For homes that are 10 doors apart, we divide not by 40 but by 36.07.

Numerical Simulations

All simulations were implemented using Matlab and run on a 100×100 house uniform lattice with no-flux boundary conditions on both the burglars and dynamic attractiveness. Having no-flux boundary conditions simply means that there is effectively a wall around the domain through which neither dynamic attractiveness nor burglars can escape. For the results that follow, the parameter values used are shown in Table 1.

Random Burglaries

Before looking at the boost and flag models, we examined the results of a model in which burglary is the result of a purely random Poisson process (hereafter the null model). It is important to do so to determine to what extent (if any) observed patterns in the subsequent models are a product of the general form of the model; that is, a Poisson process with burglars that have a finite lifetime.

For each run of this simulation, patterns are simulated over a 10-year virtual time period with static attractiveness $\tilde{A}_s = 0.0153$ (the same for every s).⁵ There is no dynamic component of attractiveness (B_s^t is set to zero for all t and s). Hence, the starting locations of newly generated burglars follow a uniform distribution across all lattice sites. In each time interval, burglars burgle with probability:

$$p = 1 - \exp\left(-\tilde{A}_s \delta t\right) \quad (8)$$

Since $\tilde{A}_s$ is constant across the domain, p is constant for all space (as well as constant in time).

Burglars perform a random walk on the lattice until they have been in the lattice for f days, at which point they are removed. Twenty-five runs of the simulation (over a virtual period of 10 years) were performed and the mean results inspected. It was clear that for the null model few homes were revictimized (none were victimized more than four times), and there was no evidence of space-time clustering. To illustrate, while an average of 2,017 burglaries occurred over the 10-year simulated interval, on average only 1.2 repeats occurred within 1 week of a previous burglary. Of course, if we increase $\tilde{A}_s$ the number of (swift) repeat victimizations would increase, but so too would the total number of burglaries. In short, for this model the frequency of repeat and near-repeat victimization (and their time course) was in line with expectation but not with the patterns observed in real-life burglary offenses.

Flag Models

Having found that the results generated by the null model were in line with the expectation, we examined a series of models to examine the flag hypothesis. To do this, we imposed on the domain a spatially inhomogeneous static component of attractiveness, $\tilde{A}_s$. Everything in the simulation is the same as that used to test the null model except $\tilde{A}_s$ is no longer constant over the domain. As before, B_s^t is set to zero for the duration of the simulation, and so does not evolve according to equation (2). This means that as with the null model there is no boost to attractiveness as a result of a burglary.

In addition to target attractiveness varying across homes, as discussed above the spatial distribution of target attractiveness may vary in systematic ways (in what follows we will use the terms risk and target attractiveness interchangeably). Consider that spatial variation may be completely random with high-risk homes neighboring high- and low-risk ones with equal likelihood. Alternatively, the variation in risk may be spatially patterned with homes closest to each other being more likely to experience similar risks than those located some distance apart. Research (for a recent discussion, see Weisburd, Bruinsma, and Bernasco 2008) suggests that crime risk is clustered in space and so complete random variation seems unlikely. However, even where target attractiveness is autocorrelated, different spatial configurations are plausible. Thus, rather than examining the role of risk heterogeneity alone, we examine how three different types of spatial configuration influence simulated patterns of crime.

Table 2. Homes victimized one to n times for each model

Number of Times Victimized (n)	Mean Number of Homes Burgled n Times (2 d.p)					
	Null	Spots	Stripes	Smoothed	Boost	Spots + Boost
1	1,645.16	851.52	639.00	1,607.52	1,566.72	760.16
2	166.08	92.44	283.04	186.72	184.72	95.68
3	12.24	38.92	127.64	15.84	16.24	39.28
4	0.64	28.12	50.52	1.76	1.68	23.64
5	0	20.08	17.60	0.16	0.08	20.80
6	0	18.28	6.44	0	0	17.88
7	0	14.32	1.56	0	0	13.32
8	0	11.44	0.28	0	0	10.72
9	0	10.56	0.04	0	0	9.80
10	0	7.56	0	0	0	8.72

The configurations used are only conceptual representations of possible distributions, but their use allows us to examine the influence of risk heterogeneity in a more systematic way. For the first configuration, we use a stripe pattern, analogous to target attractiveness varying along streets, with some streets experiencing particularly high risk. In the second, we model a scenario in which four areas are essentially crime hot spots, with risk being the greatest at the center of each spot and decaying as a function of the distance from it. In the final model, the risk allocated to each home is the function of a uniform random number generator and a smoothing function. In this model then, the risk of victimization varies across homes but the smoothing function ensures that homes that are near to each other will have some degree of similarity. Thus, for the latter model the degree of spatial autocorrelation is considerably weaker and less systematic than for the other two but is still apparent. The mean values of $\bar{A}_s$ over the domain to two significant digits are 0.0087 ($SD = 0.010$) for the stripes configuration; 0.012 ($SD = 0.0084$) for the spots configuration; and 0.015 ($SD = 0.0027$) for the final configuration.

As before, results were averaged for 25 (10-year) simulation runs for each configuration. Results are shown in Table 2, and Figures 3–5. Over the 10-year simulation period, there were an average of 1,930 burglaries for configuration 1 (stripes), 2,083 for configuration 2 (spots), and 2,036 for configuration 3 (random and smoothed). Considering the patterns of repeat victimization proper, it is evident that the patterns vary across the three configurations.




Figure 3. Results of 25 simulations for the model with heterogeneous risks across homes (spots configuration; error bars show minimum and maximum values).

With respect to the spatial patterns, for two of the configurations tested, there is evidence of spatial autocorrelation whereby neighbors of burgled homes are more likely to be victimized than those located further away. For the third, there was no pattern. There was no evidence of a temporal pattern for any of the models tested. That is, the results show that burglary is approximately as likely to be close in space to a previous burglary if that previous burglary occurred 1 or 2 weeks before. The fact that this finding is stable across the three configurations tested leads us to believe that it is unlikely that risk heterogeneity alone can explain the combined spatial and temporal patterns seen in real burglary data.

Boost Models

To examine the boost hypothesis, we examine two models. For the first, there is no risk heterogeneity, instead $\tilde{A}_s$ was given the constant value of 0.0145 for each lattice site in the domain, and the simulation is initialized with starting values for B'_s of 0.0025 across all locations. The results of 25 averaged simulation runs over a simulation time period of 10 years are shown in Figure 6. For this model, there were an average of 1,992 burglaries and more repeat victimizations than for the null model. There is also some evidence of spatial autocorrelation in the distribution of simulated




Figure 4. Results of 25 simulations for the model with heterogeneous risks across homes (stripe configuration; error bars show minimum and maximum values).




Figure 5. Results of 25 simulations for the model with heterogeneous risks across homes (smoothed configuration; error bars show minimum and maximum values).




Figure 6. Results of 25 simulations for the boost model with homogeneous risks across homes (error bars show minimum and maximum values).

burglaries. Moreover, we see that the risk of repeat and near-repeat victimization decays from the first week after a burglary event to the second. Thus, it appears that a very simple implementation of the boost hypothesis generates spatially and temporally correlated simulated burglary data under the assumptions of the model. However, the effect is subtle.

For the second model, we model the effects of both risk heterogeneity and event dependency. For this model, we use the spots configuration with a mean value of 0.0106 ($SD = 0.0077$) for $\tilde{A}_s$. This generates an average of 2,109 burglary events per simulation. Figure 7 shows that for this model, the pattern of repeat victimization and spatial autocorrelation is accentuated, and that the space–time pattern generated provides better qualitative agreement with patterns observed in real burglary data than the boost model for which risk is homogenous across homes.

Parameter values other than those shown in Table 1 could and were used, but the results generated were generally consistent with those presented above. Where alternative parameter values were used for the flag models, it was evident that none of those tested generated events that clustered in space *and* time. With respect to the boost models, increasing the values of the parameters η and ω generated results that approached those of the flag models but this is, of course, to be expected.




Figure 7. Results of 25 simulations for the boost model with heterogeneous risks across homes (spots configuration; error bars show minimum and maximum values).

Discussion

Theories of spatial crime patterns have developed considerably over the last three decades with those factors that influence crime placement being explored at ever more precise levels of resolution (i.e., places) both theoretically and empirically. However, while the importance of time is implicit in most theories, and while research has considered how crime patterns vary by time of day, with a few exceptions (e.g., Braga, Papachristos, and Hureau 2010; Spelman 1995; Weisburd et al. 2004), relatively less attention has been given to how crime patterns evolve over time, where time refers to days, weeks, or months. This is evidenced in the types of statistical analysis typically employed in research studies. For example, those with an interest in spatial patterns tend to use statistical models that focus on spatial distributions of crime, analyzing data aggregated for long intervals of time. Those with an interest in temporal patterns instead use a variety of time-series methods, but rarely are the two types of analysis integrated. Thus, for one type of analysis, time is typically ignored whereas for the other space is neglected. Much may be learned by considering the two dimensions together.

In addition to the focus on time and space, it also is important to focus on the finest level of resolution possible, in our analyses—places rather than

spaces. In the present analysis, we have focused on patterns of crime at the level of the individual household but also consider how variation in risk in the surrounding space might influence patterns of victimization.

In particular, our aim was to consider space–time patterns of burglary and to examine two theoretical models that have been proposed to explain them. The use of mathematical models allowed us to formally specify and explore model outcomes using stochastic simulations. In the introduction to this article, we discussed the possibility that observed space–time patterns of crime may be explained by a statistical artifact that occurs when results are aggregated for populations with very different risks (the flag account). However, the results of a series of simulations suggest that such models (as specified here) were insufficient and did not generate the types of pattern that are observed in real-world data. Thus, while the role of risk heterogeneity is undisputed in the generation of spatial patterns of crime, it seems unlikely that the very distinct patterns of space–time clustering that have so far proven to be ubiquitous across studies can be explained in terms of the flag explanation alone.

In contrast to the flag hypothesis, our results suggest that the boost account may offer a plausible explanation for why crime clusters in space *and* time. Of course, in reality, and as shown by our combined model, both explanations are likely to have a part to play. This seems sensible given that decades of research (e.g., Rengert and Wasilchick 2000) demonstrate the role of target attractiveness in offender decision making and that an offender has to first select a home (and they must do so using some selection criteria) before the risk to that location can be boosted. One of the challenges for future research will be to try to quantify what the relative contributions of these two mechanisms are and, if the balance between them is time stable, and whether it varies across space (and space–time) and offenders.

However, it is important to acknowledge that while the combined model did generate the qualitative patterns we sought to simulate, this does not demonstrate that it is necessary or that it is the only model that could generate them (see Eck and Liu 2008a). A range of alternative models may exist, and these may generate outcomes that are still more consistent with those of interest. Such models may involve subtle variations, such as using a multiplicative boost function (see Johnson 2008), or they may reflect very different theoretical perspectives. Without testing different models, it is not possible to evaluate whether the model implemented does explain observed patterns just that it remains a candidate explanation. Considered in concert with the cumulative findings of research that has used very different

methods, our results add further credibility to the boost hypothesis, but they do not and cannot show that it is necessary to generate the patterns of interest.

Moreover, the boost hypothesis would benefit from further attention, theoretical and empirical. In the current research and that presented elsewhere (e.g., Johnson et al. 2009), the assumption is that when a home is burgled the risk to that home and those nearby is temporarily elevated. And, that the reason for this is that the offender who committed the prior offense/offenses will return to those places for which the rewards outweigh the associated risk and effort expended. This represents a foraging model (Bernasco 2008; Johnson and Bowers 2004; Johnson et al. 2009) and assumes that the offender is a rational agent (Cornish and Clarke 1986). However, while analyses of crimes detected by the police certainly suggest that (near) repeats are typically the work of returning offenders (Bernasco 2008; Johnson et al. 2009) what remains unclear is precisely why offenders decide to return to some places but not others, and what types of offenders operate in this way. Considering the first point, it is certainly not the case that all homes are repeatedly victimized or that near repeats always follow a previous offense. Thus, determining whether there are regularities in the types of places that are most likely to encourage space-time clustering would be a logical next step. Such conditions might include particular characteristics of victimized homes or those on a block face, or other localized features of the urban backcloth. For example, near repeats may be more (or less) likely on streets segments which offer good visibility (Bowers and Johnson 2005), or on particular types of street segment (e.g., arterial or local roads). Some time ago, such research would have been unthinkable, but with the increasing availability of data at such small units of analysis and the proliferation of geographic information systems it is now a possibility.

Considering the offender as forager hypothesis, it is unlikely that all offenders adopt foraging strategies, or that they would do so all of the time. Thus, exploring the characteristics of those offenders who do adopt them and the extent to which this is their preferred strategy will be important. It may be the case that for many offenders, the use of foraging strategies is episodic or even random, whereas for others it is the dominant strategy employed. Drawing upon the ecology literature (e.g., Pyke 1984), for which theories of foraging are well established, is likely to prove valuable. The types of questions to be explored may be addressed using a variety of research methods including mathematical models, offender interviews (see Summers, Johnson, and Rengert 2010), and the analysis of crimes detected

by the police. In fact, it is unlikely that an accurate picture will be reached through the use of one method alone and so the use of different methods is encouraged.

In the current article, we have focused on the crime of burglary. However, the available evidence suggests that the type of space–time clustering we sought to explain is evident for other types of crime including vehicle-related offenses (Johnson et al. 2009), robbery and assault (Grubestic and Mack 2008), gun crime (Ratcliffe and Rengert, 2008), and even insurgent activity in Iraq (Johnson and Braithwaite, 2009; Townsley, Johnson, and Ratcliffe 2008). Thus, further research might usefully investigate the role of time-stable and dynamic factors in the generation of these crime types. One challenge for such research will involve the conceptualization of the appropriate level of place at which time-stable and dynamic factors might be modeled.

To conclude, analyses conducted at the micro level using a range of different methods suggest that there is a flux to crime that cannot be explained in terms of time-stable variation in risk across places alone, but that time-stable factors do have a part to play. One challenge for research concerned with spatial patterns of crime at the micro level then is to try to better understand not only how features of the environment influence crime risk but also how stable such factors are and at what levels (place or space) influences are exerted and experienced.

Acknowledgments

We would like to thank the reviewers and editors for their thorough review of this article and their thoughtful suggestions on how to improve it. We would also like to thank Prof. John Ockendon for useful discussions about modeling criminality and victimization and his continued encouragement.

Declaration of Conflicting Interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors disclosed receipt of the following financial support for the research and/or authorship of this article: Ashley B. Pitcher is currently supported by the “DyXi” project of the SYSCOMM program of the French National Agency for Research (grant number ANR-08-SYSC-008).

Notes

1. Patterns that are believed to describe the general distribution of real-world phenomena. For example, the time course of repeat victimization has so far proven to be ubiquitous and so can be considered a statistical signature.
2. Short et al. (2008) keep the equation deterministic by considering the expected number of burglaries instead of the random variable X_s^t .
3. We acknowledge that in the real world burglars will differ in the strategies used and some may spend much time assessing each potential target (e.g., Rengert and Wasilchick 2000). However, in the absence of data that can be used to estimate the relevant distributions, in the current article we use the same value across agents.
4. A short note on this formula, and an illustration of the edge effect for the type of analysis presented is available from the authors upon request.
5. This generates a rate of approximately 2 burglaries per 100 homes which, at the time of writing approximates the mean risk of burglary with entry in the United Kingdom (Moley 2009).

References

- Ashton, J., I. Brown, B. Senior, and K. Pease. 1998. "Repeat Victimization: Offender Accounts." *International Journal of Risk, Security and Crime Prevention* 3:269-79.
- Beavon, D. J., P. Brantingham, and P. Brantingham, P. 1994. "The Influence of Street Networks on the Patterning of Property Offenses." In *Crime Prevention Studies Vol II*, 115-148. Edited by R. Clarke. New York, NY: Willow Tree Press.
- Bernasco, W. 2008. "Them Again? Same Offender Involvement in Repeat and Near Repeat Burglaries." *European Journal of Criminology* 5:411-31.
- Bevis, C. and J. B. Nutter. 1977. "Changing Street Layouts to Reduce Residential Burglary." Paper presented to the American Society of Criminology annual meeting in Atlanta.
- Boots, B. and A. Getis. 1988. *Point Pattern Analysis*. Beverly Hills, CA: Sage.
- Bowers, K. J. and S. D. Johnson. 2005. "Domestic Burglary Repeats and Space-Time Clusters: The Dimensions of Risk." *European Journal of Criminology* 2:67-92.
- Bowers, K. J., S. D. Johnson, and K. Pease. 2005. "Victimisation and Re-Victimisation Risk, Housing Type and Area: A Study of Interactions." *Crime Prevention and Community Safety* 7:7-17.
- Budd, T. 1999. *Burglary of Domestic Dwellings: Findings From the British Crime Survey*. Home Office Statistical Bulletin, 4/99. London, England: Home Office.
- Braga, A., A. V. Papachristos, and D. M. Hureau. 2010. "The Concentration and Stability of Gun Violence at Micro Places in Boston, 1980-2008." *Journal of Quantitative Criminology* 26:33-53.

- Brantingham, P. and P. Brantingham. 1993. "Environment, Routine, and Situation: Toward a Pattern Theory of Crime." Pp. 259-94 in *Routine Activity and Rational Choice*, edited by R. V. Clarke and M. Felson. New Brunswick, NJ: Transaction Publishers.
- Brantingham, P. and P. Brantingham. 2008. "Crime Pattern Theory." Pp. 78-93 in *Environmental Criminology and Crime Analysis*, edited by R. Wortley and L. Mazerolle. Portland, OR: Willan.
- Cornish, D. and R. Clarke. 1986. *The Reasoning Criminal: Rational Choice Perspectives on Offending*. New York, NY: Springer-Verlag.
- Eck, J. and L. Liu. 2008a. "Contrasting Simulated and Empirical Experiments in Crime Prevention." *Journal of Experimental Criminology* 4:195-213.
- Eck, J. and L. Liu. 2008b. "Varieties of Artificial Crime Analysis: Purpose, Structure, and Evidence in Crime Simulations." Pp. 413-32 in *Artificial Crime Analysis Systems: Using Computer Simulations and Geographic Information Systems*, edited by L. Liu and J. Eck. Hershey, PA: Idea Group PLC.
- Eck, J. and D. Weisburd. 1995. "Crime Places in Crime Theory." Pp. 3-31 in *Crime and Place*, edited by J. Eck and D. Weisburd. New York, NY: Criminal Justice Press.
- Ericsson, U. 1995. "Straight from the Horse's Mouth." *Forensic Update* 43:23-5.
- Farrell, G. 2005. "Progress and Prospects in the Prevention of Repeat Victimization." Pp. 145-172 in *Handbook of Crime Prevention and Community Safety*, edited by N. Tilley. Cullompton, UK: Willan.
- Gilbert, N. 2008. *Agent-Based Models*. Thousand Oaks, CA: Sage.
- Grubestic, T. H. and E. A. Mack. 2008. "Spatio-Temporal Interaction of Urban Crime." *Journal of Quantitative Criminology* 24:285-306.
- Hillier, B. 2004. "Can Streets be Made Safer?" *Urban Design International* 9:31-45.
- Johnson, S. D. 2008. "Repeat Burglary Victimization: A Tale of Two Theories." *Journal of Experimental Criminology* 4:215-40.
- Johnson, S. D., Bernasco, W., Bowers, K. J., Elffers, H., Ratcliffe, J., Rengert, G., Townsley, M. 2007. "Near Repeats: A Cross National Assessment of Residential Burglary." *Journal of Quantitative Criminology* 23:201-19.
- Johnson, S. D. and K. J. Bowers. 2004. "The Stability of Space-Time Clusters of Burglary." *British Journal of Criminology* 44:55-65.
- Johnson, S. D., Braithwaite, A. 2009. "Spatio-Temporal Distribution of Insurgency in Iraq." In J. Freilich & G. Newman (eds.), 'Countering Terrorism through SCP.' *Crime Prevention Studies* 25:9-32.
- Johnson, S. D. and K. J. Bowers. 2010. "Permeability and Burglary Risk: Are Cul-de-Sacs Safer?" *Journal of Quantitative Criminology* 26:89-111.
- Johnson, S. D., K. Bowers, and A. Hirschfield. 1997. "New Insights in the Spatial and Temporal Distribution of Repeat Victimization." *British Journal of Criminology* 37:224-41.

- Johnson, S. D., L. Summers, and K. Pease. 2009. "Offenders as Forager: A Direct Test of the Boost Account of Victimization." *Journal of Quantitative Criminology* 25:181-200.
- Kleemans, E. R. 2001. "Repeat Burglary Victimization: Results of Empirical Research in the Netherlands. Pp. 53-68 in *Crime Prevention Studies 12*, edited by G. Farrell and K. Pease. Monsey: CRC press.
- Moley, S. (2009). "Property Crime." Pp. 73-87 in *Crime in England and Wales 2008/09*, edited by A. Walker, J. Flatley, C. Kershaw, and D. Moon. London, England: Home Office.
- Morgan, F. 2001. "Repeat Burglary in a Perth Suburb: Indicator of Short-Term or Long-Term Risk?" Pp. 83-118 in *Repeat Victimization: Crime Prevention Studies*. Vol. 12, edited by G. F. Pease. Monsey: Criminal Justice Press.
- Nelson, J. 1980. "Multiple Victimization in American Cities: A Statistical Analysis of Rare Events." *American Journal of Sociology* 85:870-91.
- Osborn, D. R., and Tseloni, A. 1998. The distribution of household property crimes. *Journal of Quantitative Criminology*, 14:307-330.
- Pease, K. 1998. *Repeat Victimization: Taking Stock*. Home Office Police Research Group, Crime Detection and Prevention Series, Paper 90. London, England: Home Office.
- Pyke, G. 1984. "Optimal Foraging Theory: A Critical Review." *Annual Review of Ecology and Systematics* 15:523-75.
- Polvi, N., T. Looman, C. Humphries, and K. Pease. 1991. "The Time Course of Repeat Burglary Victimization." *British Journal of Criminology* 31:411-14.
- Ratcliffe, J. H. 2004. "The Hotspot Matrix: A Framework for the Spatio-Temporal Targeting of Crime Reduction." *Police Practice and Research* 5:5-23.
- Ratcliffe, J. H. and Rengert, G. F. 2008. "Near Repeat Patterns in Philadelphia Shootings." *Security Journal*, 21:58-76.
- Rengert, G. F. and J. Wasilchick. 2000. *Suburban Burglary: A Tale of Two Suburbs*. 2nd ed. Springfield, IL: Charles C. Thomas.
- Robinson, W. 1950. "Ecological Correlations and the Behavior of Individuals." *American Sociological Review* 15:351-7.
- Shaw, C. and H. McKay. 1969. *Juvenile Delinquency and Urban Areas*. Chicago, IL: University of Chicago Press.
- Sherman, L., P. Gartin, and M. Buerger. 1989. "Hot Spots of Predatory Crime: Routine Activities and the Criminology of Place." *Criminology* 27:27-55.
- Shipley, R. J. and S. J. Chapman. 2010. "Multiscale Modelling of Fluid and Drug Transport in Vascular." *Bulletin of Mathematical Biology* 72:1464-91. doi:10.1007/s11538-010-9504-9.

- Short, M., P. J. Brantingham, A. L. Bertozzi, and G. E. Tita. 2010. "Dissipation and Displacement of Hotspots in Reaction-Diffusion Models of Crime." *Proceedings of the National Academy of Sciences of the United States* 107:3961-5.
- Short, M., M. R. D'Orsogna, P. J. Brantingham, and G. E. Tita. 2009. "Measuring and Modelling Repeat and Near Repeat Effects." *Journal of Quantitative Criminology* 25:325-9.
- Short, M., M. R. D'Orsogna, V. B. Pasour, G. E. Tita, P. Brantingham, A. L. Bertozzi, and L. B. Chayes. 2008. "A Statistical Model of Criminal Behavior." *Mathematical Models and Methods in Applied Sciences* 18:1249-67.
- Sparks, R. 1981. "Multiple Victimisation: Evidence, Theory and Future Research." *Journal of Criminal Law and Criminology* 72:762-78.
- Spelman, W. 1995. "Criminal Careers of Public Places." Pp. 115-44 in *Crime and Place*, edited by J. Eck and D. Weisburd. New York: Criminal Justice Press.
- Summers, L., S. D. Johnson, and G. Rengert. 2010. "The Use of Maps in Offender Interviews." Pp. 246-72 in *Offenders on Offending: Learning About Crime From Criminals*, edited by W. Bernasco. Cullompton, UK: Willan.
- Townsley, M., R. Homel, and J. Chaseling. 2000. "Repeat Burglary Victimization: Spatial and Temporal Patterns." *Australian and New Zealand Journal of Criminology* 33:37-63.
- Townsley, M., R. Homel, and J. Chaseling. 2003. "Infectious Burglaries: A Test of the Near Repeat Hypothesis." *British Journal of Criminology* 43:615-33.
- Townsley, M. T., S. D. Johnson, and J. R. Ratcliffe. 2008. "Space-Time Dynamics of Insurgent Activity in Iraq." *Security Journal* 21:139-46.
- Vaupel, J. W. and A. I. Yashin. 1985. "Heterogeneity's Ruses: Some Surprising Effects on Selection on Population Dynamics". *The American Statistician* 39:176-85.
- Weisburd, D., G. J. Bruinsma, and W. Bernasco. 2008. "Units of Analysis in Geographic Criminology: Historical Development, Critical Issues and Open Questions." Pp. 3-34 in *Putting Crime in its Place: Units of Analysis in Geographic Criminology*, edited by D. Weisburd, W. Bernasco, and G. J. Bruinsma. New York: Springer.
- Weisburd, D., S. Bushway, C. Lum, and S. Yang. 2004. "Trajectories of Crime at Places: A Longitudinal Study of Street Segments in the City of Seattle." *Criminology* 42:283-321.
- Xie, M., G. Pogarsky, J. P. Lynch, and D. McDowall. 2006. "Prior Police Contact and Subsequent Victim Reporting: Results From the NCVS." *Justice Quarterly* 23:481-501.

Bios

Ashley B. Pitcher is a postdoctoral researcher employed by the French National Center for Scientific Research (CNRS) and is hosted by the Centre d'Analyse et de Mathématique Sociales (CAMS) at the School for Advanced Studies in the Social Sciences (EHESS). She received her doctorate in mathematics from the University of Oxford, in 2009. While at Oxford, she was a member of the Oxford Centre for Industrial and Applied Mathematics (OCIAM) and Balliol College. Her research interests include applications of mathematics in the social sciences.

Shane D. Johnson is a reader at the UCL Department of Security and Crime Science, UCL Jill Dando Institute of Crime Science, University College London, England. He has a PhD and an MA in psychology and a BSc in computer science. He has particular interests in the spatial and temporal distribution of crime, computer simulation, and evaluation methods.