



What Do Women Want? Looking Beyond Patient Satisfaction

Deborah McCarter & Carrie E. MacLeod

ABSTRACT

Objective: To hear the voices of women, their partners, and nurses about expectations and priorities during the postpartum hospitalization.

Design: Focus groups using semistructured interview questions.

Setting: A 12-bed labor–delivery–recovery–postpartum unit at a small urban hospital in the U.S. Northeast.

Participants: Women who planned to or had given birth, their partners, and the maternity nurses who cared for them.

Measurements: Qualitative thematic analysis of focus group transcripts.

Results: Thematic analysis produced the following themes for women's priorities: *Need for individualized attention to maternal physical and emotional care; Fear of providing inadequate care for the newborn, including establishing infant feeding; and Transitioning to parenting as a new mother versus as an experienced mother.* Themes for nurses' priorities included *Safety issues around sleep and breastfeeding, Transitioning to*

parenting with an emphasis on maternal self-care, and Addressing barriers to effective discharge education. Response comparisons between the women/partners and nurses suggest that there is a disconnection between women's and nurses' priorities and expectations for care during the postpartum period.

Conclusion: Women and nurses identified unmet needs in the postpartum period, consistent with the current literature. Providing standardized education during the transitional period around discharge from the hospital to home may not be optimal and may even detract from meeting the needs for rest and connection with family and the health care team. Nursing care that extends beyond the maternity hospitalization may be needed to individualize care and meet previously unmet needs.

doi: [10.1016/j.nwh.2019.09.002](https://doi.org/10.1016/j.nwh.2019.09.002)

Accepted September 2019

KEYWORDS: discharge teaching, fourth trimester, health education, maternal satisfaction, patient satisfaction, postpartum

CLINICAL IMPLICATIONS

- Maternity care during the postpartum hospitalization blends responsibility for ensuring physical recovery for women, supporting newborns' transition to extrauterine life, establishing newborn feeding, and educating families on newborn care and safety.
- For maternal–child nurses working in the postpartum setting, the importance of meeting the many needs of families creates an urgency around postpartum discharge education.
- There is often a disconnect between women's and nurses' priorities during this time; such conflicting priorities can contribute to decreased satisfaction among women receiving care.
- Standardized discharge education may not be effective in meeting the needs of women and their families.
- Nursing expertise could be used to extend care beyond discharge.

Maternal–newborn care, at its best, is focused on health promotion during a critical developmental phase for families. To that end, education is a key aspect of the nursing role in maternity care. In fact, education is a deeply entrenched and valued professional nursing function (Bastable, 2008). Maternity care during the postpartum hospitalization blends responsibility for ensuring physical recovery for women, supporting newborns' transition to extrauterine life, establishing newborn feeding, and educating families on newborn care and safety. There is much to accomplish during this brief hospitalization, and this is conducted in an environment with many visitors, unpredictable newborn demands, and, often, fatigued mothers.

Reducing risks for maternal mortality and morbidity has become a public health priority (Centers for Disease Control and Prevention, 2017a, 2017b). Guidelines and tools for nurses to help address this need have been developed by the Association of Women's Health, Obstetric and Neonatal Nurses (AWHONN; n.d.), with the goal of reducing postpartum complications through focused education. In addition, since 2016, a multidisciplinary initiative known as the 4th Trimester Project (University of North Carolina Center for Maternal and Infant Health, n.d.) has raised awareness of the many unmet physical and psychosocial needs identified by women and their health care providers (Lu et al., 2006; Martin, Horowitz, Balbierz, & Howell, 2014; Tully, Stuebe, & Verbiest, 2017).

Deborah McCarter, PhD, RN, IBCLC, WHNP-BC, is a professor, Saint Anselm College, Manchester, NH. **Carrie E. MacLeod**, PhD, APRN-BC, is an assistant professor, Saint Anselm College, Manchester, NH. The authors report no conflicts of interest or relevant financial relationships. **Address correspondence to:** dmccarter@anselm.edu.



Women identified many distractors and stressors during the first few days postpartum, which made it even more difficult to remember information that was conveyed during discharge teaching

For maternal–child nurses working in the postpartum setting, the importance of meeting the many needs of families creates an urgency around postpartum discharge education. Accrediting bodies such as The Joint Commission rightly seek documentation that patient safety goals are being met (The Joint Commission, 2019). Initiatives to improve care based on patient feedback, such as Press Ganey (2019), focus on goals of patient satisfaction. Professional organizations highlight the need for better preparation, recognition, and response to complications such as postpartum hemorrhage (AWHONN, 2017). Nurses juggle all these priorities in a fast-paced and frequently changing environment. It is no wonder that many standardized checklists are created to ensure that no topics are missed before discharge.

Although nurses and other health care providers are charged with providing ever more content, there is little evidence to determine whether the breadth of discharge teaching is received and retained by women and families in this transitional life change, and there is some evidence that it is not (McCarter-Spaulling & Shea, 2016). Despite the large proportion of valuable nursing time dedicated to discharge teaching, current systems appear to be inadequate to address postpartum medical and psychosocial risks (Lu et al., 2006) or to meet women's identified needs for support with breastfeeding, newborn care, maternal self-care, and postpartum depression (Kanotra et al., 2007; Kleppel, Suplee, Stuebe, & Bingham, 2016; Tully et al., 2017). Standardization of the topics to be addressed may compete with individualized care based on maternal/newborn/family needs and priorities.

TABLE 1 FOCUS GROUP PARTICIPANTS

Population Sampled	Participants, <i>n</i>	Participant Characteristics
Women who recently participated in a research study measuring outcomes of education about postpartum depression	9	Six women after childbirth with between two and six children at home One woman pregnant with second child Two pregnant women, each with one child at home
Women presenting for prenatal care	2	One adolescent couple expecting their first child
Participants of postpartum mother–baby groups	4	Two first-time mothers and two mothers with second children, all with newborns
Women presenting for pediatric or postpartum care or childbirth classes	2	One woman and her partner who recently had their third child
	3	One first-time mother with newborn One first-time mother whose child was now 7 years old One woman currently pregnant with her third child
	4	One couple pregnant with first child One first-time mother with a toddler One woman with other children and a newborn

Our purpose was to listen to the voices of women, their partners, and nurses about their expectations and priorities for education and anticipatory guidance during the postpartum hospitalization. These results are being used to inform strategies being developed by the hospital's postpartum education committee to improve nurses' ability to provide individualized care that also prioritizes safety.

Design

For members of the postpartum education committee to understand more clearly what women and families expected and desired, we convened five focus groups for women and their partners and two focus groups for nurses. Our aim was to provide insight into whether nurses and childbearing women identified the same priorities for postpartum care. Currently, in this hospital, women are given a standardized list of content areas to be covered, and they indicate which areas they would like to address with their nurse. This content is also available in a written booklet. During the hospitalization, nurses address each of these content areas with a woman and her family if desired. Once these items are addressed, the nurse and the woman sign paperwork indicating that teaching has been completed. This form must be completed before discharge.

Sample

We recruited participants for this convenience sample from women who had given birth or planned to give birth at a 12-bed maternity unit of a small urban hospital in the U.S. Northeast, at which there are approximately 1,000 births/year, and from the nursing staff of the hospital. After approval by the hospital's institutional review board, we convened a total of seven focus groups between 2015 and 2017—five with women/

partners and two with nurses. We offered focus group participation to recent participants in a research study of postpartum discharge education, and women/partners who presented for care at prenatal, postpartum, or childbirth education settings (see Table 1). A total of 24 parents, representing the prenatal and postpartum periods, participated in focus groups. We attempted to have separate groups for prenatal and postpartum respondents, but scheduling for interested respondents necessitated holding groups when the most people were available. The number of groups that could be held was limited by the availability of funding, so data saturation was likely not attained. We offered participation in the nurses' groups to any staff nurse from the labor–delivery–recovery–postpartum unit and the Level 2 special care nursery and ultimately included a total of 16 nurses between the two groups.

We held all groups at the maternity hospital; each lasted 2 hours and included a meal. We provided participants in the women/partners groups with a \$25 cash incentive after the group meeting. Nurses received documentation in support of their professional development. All participants gave written informed consent at the beginning of the meeting and agreed to be recorded. Research assistants participated by assisting with greeting, ensuring comfort, observing group dynamics, and providing the payments at the end of the group.

We conducted semistructured interviews using a predetermined interview question guide. The semistructured format provided a balance between an interviewer working flexibly with the interview guide and permitting time for participants' individualized narratives. We asked open-ended questions to enable participants to provide responses in their own words. If necessary, we used prompting questions to further expand on participants' narratives. We recorded the

groups with a digital audio recorder, and the recordings were transcribed professionally into written form for analysis.

We used the six-phase thematic analysis recommended by Braun and Clarke (2006). The six phases are (a) familiarization with data, (b) generation of initial codes, (c) search for themes, (d) review of themes, (e) definition and naming of themes, and (f) production of report. We also used strategies outlined by Lincoln and Guba (1985) for transferability, dependability, confirmability, and credibility to establish research trustworthiness (Krefting, 1991).

Results

Thematic analysis provided emergent themes related to women's and nurses' perceived priorities around discharge teaching. Themes for women's priorities included *Need for individualized attention to maternal physical and emotional care*, *Fear of providing inadequate care for the newborn, including establishing infant feeding*, and *Transitioning to parenting as a new mother versus as an experienced mother*. Themes for nurses' priorities included *Safety issues around sleep and breastfeeding*, *Transitioning to parenting with an emphasis on maternal self-care*, and *Addressing barriers to effective discharge education*.

Need for Individualized Attention to Maternal Physical and Emotional Care

Women consistently expressed that the attention paid to physiologic changes far exceeded the attention given to their emotional needs. The combination of physical changes and sleep deprivation had many wondering, "When will I feel normal again?" Women also expressed a feeling of being forgotten during the time between discharge from the hospital and their first postpartum visit. A priority for the women who participated in this study was more individualized attention that focused on emotional and physical changes.

- "I feel emotional needs and change . . . gets pushed to the side or it's neglected a little bit. . . . The focus is so much like on physiological stuff. I remember I was really blindsided. . . ."
- "And the next time you get checked out it's 6 weeks later . . . and the baby gets checked out multiple times . . . You're kind of forgotten."
- "I need you [nurse] to be an extension of my family . . . so it's as easy as it possibly can be."
- "... different reaction from me if you asked me face to face versus a questionnaire [depression screening]. I felt like if somebody was there and you had to look someone in the eye, it would be a lot more personal."

Fear of Providing Inadequate Care for the Newborn, Including Establishing Infant Feeding

Women's comments reflected a sense of catastrophic thinking regarding care of the newborn and establishing

newborn feeding. They wondered whether they were giving their newborns appropriate care and nutrition. Many shared the feeling that they "didn't know what they didn't know" or that they had forgotten essential information regarding newborn care. Women expressed a feeling of self-reproach because they felt they were providing inadequate care. Many shared a sense of failure that they did not remember everything they had learned about newborn care or that they were unable to focus as they tried to remember everything.

- "I'm afraid there will be one thing on the list that he's not doing or that I haven't noticed him doing."
- "I remember the first baby . . . didn't even know what I didn't know. I went home in 3 days sobbing."
- "Everyone says breastfeeding is the best but no one talks about how difficult it is. . . . I still don't feel like I know what I'm doing."

Transitioning to Parenting as a New Mother Versus as an Experienced Mother

There were distinctly different themes between first-time mothers and experienced mothers. First-time mothers shared that they felt as if they had no other choice but to care for their newborns. They struggled because they often felt like the child within this new experience. They expressed anxiety about leaving the hospital and being solely responsible for their newborns.

- "We've never done this before, so of course we're a child in that way."
- "I was a little bit anxious about leaving. I didn't know what I was doing. I was surprised they were letting me leave with a kid."

Experienced mothers shared that they felt more prepared for the birthing process but stressed that every pregnancy and birth are different and that they still required help. There was a call for nurses to not assume a woman was "good to go" because she was not a new mother. Experienced mothers' concerns included the perception that they remembered everything from previous births, breastfeeding issues, and how to reconcile the desire to return home to their other children with being petrified to "take it all on."

- "I was a little bit less wrapped up in my brain the second time around."
- "My second one—I was thinking, 'Holy crap, what did I just do? I have two kids now.'"

Nurses' perceived priorities around discharge education focused on *Safety issues around sleep and breastfeeding*, *Transitioning to parenting with an emphasis on maternal self-care*, and *Addressing barriers to effective discharge education*.

Safety Issues Around Sleep and Breastfeeding

Nurses expressed that safe sleep practices were an essential part of the discharge education. They acknowledged that, in general, new mothers seemed underinformed regarding safe sleep products. In addition, cultural and familial traditions

often superseded the advice given during discharge education regarding safe sleep and breastfeeding.

- “There are a lot of products on the market that babies sleep really well in that are really unsafe.”
- “It’s what Grandma is telling them, what culture does.”
- “I feel like a lot of women unless you’re really committed to breastfeeding and you know how to resist your mother and mother-in-law who are just telling you, ‘Just give them a bottle, it will be okay.’ That’s harder than anything.”
- “How to do . . . not battle, but counteract years of acculturation and this is what is best for your baby.”
- “. . . many people are on the verge of neglectful about really knowing how often a baby needs to eat and they can be told over and over again and they’re just not taking it in.”

Transitioning to Parenting With an Emphasis on Maternal Self-Care

Nurses appeared to agree that when discharged to home, all mothers, new and experienced, are faced with many challenges. Nurses recognized that many mothers meet these challenges by sacrificing focus on self and self-care. Nurses conceded the limited impact that postpartum discharge education may have on these individuals and stressed the need for ongoing resources and support for new and experienced mothers.

- “. . . [mother] feeling responsible for everything except for yourself, taking care of yourself.”
- “We expect them to take care of their own bodies and their own needs and their baby’s.”
- “The really big thing is making sure that they’re aware of their resources and who to call and when to call and why.”
- “You’re thinking about how nervous they are going home and how much they don’t know and they’re becoming aware of what they don’t know.”
- “First babies . . . I probably wrongly assume to give them a little more attention and probably wrongly assume with 2nd, 3rd, 4th babies . . . she’s got this.”

Addressing Barriers to Effective Discharge Education

Subthemes for this theme included woman, nurse, and system issues. Nurses’ comments reflected that women do not know what they need at the time of discharge and that new mothers are often overwhelmed by the amount of information they are given. Nurse feel challenged to find the time to be intentional and present with new mothers. This may be due to other patient assignments, staffing issues, or finding an appropriate time to converse and personalize discharge teaching.

- “I feel like it’s rare that you find a time where you can sit down with them and they don’t have any visitors and they’re receptive to information and they’re able to ask questions. I feel like that is a rare moment.”

- “We bombard them with so much stuff and they can’t take it in and then they’re so scared.”
- “I think they don’t know what they want from us.”
- “. . . but sometimes we just have to be, okay, it’s time for this to get done.”
- “Patients don’t think you’re really doing anything unless you’ve shown them a checklist and that you’ve done it.”
- “I think a big thing is teaching people that they have power over this and to empower them to feel confident to be going home and I think that’s a big part of what our teaching is supposed to be.”
- “. . . it provides a lot of nurse satisfaction when you’re actually able to have the time to sit down and personalize it and really see where those parents are coming from.”

Implications for Practice

Postpartum nurses are aware of the challenges and distractions in providing education to new families during the brief hospitalization. Women’s responses in the focus group suggest that there is a disconnection between women’s and nurses’ priorities and expectations for care during the postpartum period. It is likely that providing standardized education during this transitional period is not effective and may even detract from meeting the needs for rest and connection with family and the health care team. Differences in expectations among patients and nurses was also identified by [Tobiano, Marshall, Bucknall, and Chaboyer \(2016\)](#) in a study conducted in Australia that addressed the experience of patient-centered care on a medical unit.

Pressure to accomplish tasks such as completing discharge paperwork and documenting that all essential topics have been covered works against a nurse’s ability to provide individualized, client-centered care ([Tobiano et al., 2016](#)). These conflicting priorities have the potential to decrease women’s and nurses’ satisfaction, because communication with the nurse is a key component of perceived care ([Senti & LeMire, 2011](#)). Women’s satisfaction is considered a key element of quality, and measurement of it may even be a requirement for reimbursement. In this environment, perhaps it is time to reevaluate traditional methods for providing discharge teaching. Nurses could bring their expertise in client-centered care to clinical practice committees and address whether changes to discharge procedures and standards are needed.

Improving the experience for women and families during the maternity stay is not the only goal in teaching. Women in the focus groups described a need to have someone help them understand what is normal during the postpartum period for themselves and their infants. In a study of outcomes associated with readiness for postpartum discharge, [Bernstein et al. \(2013\)](#) found that women who were uncertain about maternal and infant symptoms and experienced concern in the first few weeks made more phone calls to health care providers (maternity and pediatric) and had lower



Similarly, our focus group participants identified many distractors and stressors during the first few days postpartum, which made it even more difficult to remember information that was conveyed during discharge teaching (Russell, 2006; Schwabe & Wolf, 2010). Perhaps increased access to health care providers after discharge is desirable, which could free up more time during the hospitalization for rest, bonding, and nurturing. Without multiple tasks to accomplish, nurses could provide more individualized care, improving satisfaction and using nursing time more effectively.

Access to continued nursing care after discharge, when women and their partners have increased readiness and ability to learn new information, could be a preferable option instead of more intensive teaching during the hospital stay

scores on measures of physical and mental health. Other variables also influenced outcomes independently, including cesarean birth, breastfeeding, poorer maternal physical and mental health, and inadequate health insurance (Bernstein et al., 2013). Although families are very motivated to learn during the postpartum period, the shortage of time during the maternity hospitalization, as well as stress and lack of confidence in negotiating the transition to parenthood, creates barriers to learning (Russell, 2006). This suggests that nurses may need to address these barriers before providing discharge teaching, in recognition of how little information may actually be retained.

Although increased health care use and, thus, the cost of care may be considered a negative outcome, it might motivate reconsideration of how postpartum care is provided. Bernstein et al.'s (2013) findings suggest that a short hospitalization does not allow enough time for education and support, particularly for families at greater risk for adverse outcomes.

Education provided during the hospitalization could focus on only the most essential needs and safety concerns. Access to continued nursing care after discharge, when women and their partners have increased readiness and ability to learn new information, could be a preferable option instead of more intensive teaching during the hospital stay. More research is needed to identify the most critical goals for teaching during hospitalization and the best methods to meet them.

The findings of this study are limited to small groups of mothers and partners from one hospital in the U.S. Northeast. However, there is little evidence measuring outcomes of nurse discharge teaching. There does not appear to be any standard for what outcomes are desirable or how to measure them (Barkin, Wisner, Bromberger, Beach, & Wisniewski, 2010). Surveys of women's satisfaction provide some information but are not adequate as an outcome measure of quality nursing care and teaching.

Conclusion

The results from these focus groups indicate that women and nurses feel that significant needs for postpartum care may be going unmet. Addressing the current systems and protocols for providing postpartum care and teaching during the maternity hospitalization and identifying creative ways to extend nursing care after discharge are necessary to better meet the needs of women and to effectively use nursing time. Future research to measure woman-centered

outcomes beyond maternal satisfaction is needed to support evidence-based nursing practice in the fourth trimester. **NWH**

Acknowledgement

Research supported by New Hampshire-INBRE through an Institutional Development Award (IDeA), P20GM103506, from the National Institute of General Medical Sciences of the NIH. The authors acknowledge the contribution of the maternity nurses and childbearing women at Catholic Medical Center, who inspire excellence in nursing care and research.



References

- Association of Women's Health, Obstetric and Neonatal Nurses. (n.d.). POST-BIRTH warning signs education program. Retrieved from <https://www.awhonn.org/page/POSTBIRTH>
- Association of Women's Health, Obstetric and Neonatal Nurses. (2017). *The AWHONN postpartum hemorrhage project*. Retrieved from <http://www.pphproject.org/resources.asp>
- Barkin, J. L., Wisner, K. L., Bromberger, J. T., Beach, S. R., & Wisniewski, S. R. (2010). Assessment of functioning in new mothers. *Journal of Women's Health*, 19(8), 1493–1499. <https://doi.org/10.1089/jwh.2009.1864>
- Bastable, S. B. (2008). *Nurse as educator: Principles of teaching and learning for nursing practice*. Boston: Jones and Bartlett Publishers.
- Bernstein, H. H., Spino, C., Lalama, C. M., Finch, S. A., Wasserman, R. C., & McCormick, M. C. (2013). Unreadiness for postpartum discharge following healthy term pregnancy: Impact on health care use and outcomes. *Academic Pediatrics*, 13(1), 27–39. <https://doi.org/10.1016/j.acap.2012.08.005>
- Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology*, 3(2), 77–101. <https://doi.org/10.1191/1478088706qp0630a>
- Centers for Disease Control and Prevention. (2017a). *Pregnancy mortality surveillance system*. Retrieved from <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pmss.html>
- Centers for Disease Control and Prevention. (2017b). *Severe maternal morbidity in the United States*. Retrieved from <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/severematernalmorbidity.html>
- Kanotra, S., D'Angelo, D., Phares, T. M., Morrow, B., Barfield, W. D., & Lansky, A. (2007). Challenges faced by new mothers in the early postpartum period: An analysis of comment data from the 2000 Pregnancy Risk Assessment Monitoring System (PRAMS) survey. *Maternal and Child Health Journal*, 11(6), 549–558. <https://doi.org/10.1007/s10995-007-0206-3>
- Kleppel, L., Suplee, P., Stuebe, A., & Bingham, D. (2016). National initiatives to improve systems for postpartum care. *Maternal and Child Health Journal*, 20, 66–70. <https://doi.org/10.1007/s10995-016-2171-1>
- Krefting, L. (1991). Rigor in qualitative research: The assessment of trustworthiness. *American Journal of Occupational Therapy*, 45(3), 214–222. <https://doi.org/10.5014/ajot.45.3.214>
- Lincoln, Y. S., & Guba, E. G. (1985). *Naturalistic inquiry*. Beverly Hills, CA: Sage Publications.
- Lu, M. C., Kotelchuck, M., Culhane, J. F., Hobel, C. J., Klerman, L. V., & Thorp, J. M., Jr. (2006). Preconception care between pregnancies: The content of prenatal care. *Maternal and Child Health Journal*, 10(Suppl. 1), S107–S122. <https://doi.org/10.1007/s10995-006-0118-7>
- Martin, A., Horowitz, C., Balbierz, A., & Howell, E. (2014). Views of women and clinicians on postpartum preparation and recovery. *Maternal and Child Health Journal*, 18(3), 707–713. <https://doi.org/10.1007/s10995-013-1297-7>
- McCarter-Spaulling, D., & Shea, S. (2016). Effectiveness of discharge education on postpartum depression. *MCN, The American Journal of Maternal/Child Nursing*, 41(3), 168–172. <https://doi.org/10.1097/NMC.0000000000000236>
- Press Ganey. (2019). *Patient experience*. Retrieved from <http://www.pressganey.com/solutions/patient-experience>
- Russell, S. S. (2006). An overview of adult-learning processes. *Urologic Nursing*, 26(5), 349–370.
- Schwabe, L., & Wolf, O. T. (2010). Learning under stress impairs memory formation. *Neurobiology of Learning and Memory*, 93(2), 183–188. <https://doi.org/10.1016/j.nlm.2009.09.009>
- Senti, J., & LeMire, S. (2011). Patient satisfaction with birthing center nursing care and factors associated with likelihood to recommend institution. *Journal of Nursing Care Quality*, 26(2), 178–185. <https://doi.org/10.1097/NCQ.0b013e3181fe93e6>
- The Joint Commission. (2019). *Hospital: 2019 national patient safety goals*. Retrieved from https://www.jointcommission.org/hap_2017_npsgs
- Tobiano, G., Marshall, A., Bucknall, T., & Chaboyer, W. (2016). Activities patients and nurses undertake to promote patient participation. *Journal of Nursing Scholarship*, 48(4), 362–370. <https://doi.org/10.1111/jnu.12219>
- Tully, K. P., Stuebe, A. M., & Verbiest, S. B. (2017). The fourth trimester: A critical transition period with unmet maternal health needs. *American Journal of Obstetrics & Gynecology*, 217(1), 37–41. <https://doi.org/10.1016/j.ajog.2017.03.032>
- University of North Carolina Center for Maternal and Infant Health. (n.d.). *4th Trimester project*. Retrieved from <http://4thtrimester.web.unc.edu>