

The influence of matrix quality on species richness in remnant forest

Ian J. Reider · Maureen A. Donnelly · James I. Watling

Received: 22 December 2017 / Accepted: 30 May 2018 / Published online: 6 June 2018
© Springer Nature B.V. 2018

Abstract

Context Habitat destruction is the leading threat to terrestrial biodiversity, isolating remnant habitat in a matrix of modified vegetation.

Objectives Our goal was to determine how species richness in several broad taxonomic groups from remnant forest was influenced by matrix quality, which we characterized by comparing plant biomass in forest and the surrounding matrix.

Methods We coupled data on species-area relationships (SARs) in forest remnants from 45 previously published studies with an index of matrix quality calculated using new estimates of plant biomass derived from satellite imagery.

Results The effect size of SARs was greatest in landscapes with low matrix quality and little forest cover. SARs were generally stronger for volant than for non-volant species. For the terrestrial taxa included in our analysis, matrix quality decreased as the

proportion of water, ice, or urbanization in a landscape increased.

Conclusions We clearly demonstrate that matrix quality plays a major role in determining patterns of species richness in remnant forest. A key implication of our work is that activities that increase matrix quality, such as active and passive habitat restoration, may be important conservation measure for maintaining and restoring biodiversity in modified landscapes.

Keywords Connectivity · Dispersal · Habitat loss · Habitat modification · Isolation · Patch

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10980-018-0664-6>) contains supplementary material, which is available to authorized users.

I. J. Reider · J. I. Watling (✉)
Department of Biology, John Carroll University,
University Heights, OH 44118, USA
e-mail: jwatling@jcu.edu

M. A. Donnelly
Department of Biological Sciences, Florida International
University, Miami, FL 33199, USA

Introduction

Habitat loss is the primary threat to terrestrial biodiversity (Barnosky et al. 2012; Newbold et al. 2016). The link between habitat loss and biodiversity maintenance is rooted in the species-area relationship (SAR), which states that large areas of habitat harbor more species than small habitat areas. Large habitats are relatively species-rich because they are better buffered from stochastic extinction events (Gonzalez and Chaneton 2002), represent bigger targets for colonists (Jones et al. 2015), and contain more resources (Steinmann et al. 2011) than small areas. The benefits of large areas for biodiversity provide the key rationale for conserving large, continuous habitat patches (Paz Durán et al. 2016), a hallmark of global

conservation efforts (Edgar et al. 2014; Watson et al. 2014).

Although habitat loss is recognized as the major threat to biodiversity, removal of habitat is usually accompanied by fragmentation, a change in landscape configuration resulting from the breaking apart of habitat (Fahrig 2003). Whereas habitat loss is a process in which the total amount of habitat in a landscape decreases, and the mean distance among patches of remnant habitat increases, fragmentation is the pattern that frequently emerges in such landscapes, wherein the number of habitat remnants and total edge length increase, and mean remnant size decreases (Fahrig 2003). Much debate has focused on the relative importance of habitat loss and fragmentation for biodiversity maintenance (Fahrig 2013; Rybicki and Hanski 2013). However, the fact that habitat loss and fragmentation are often strongly correlated (Haila 2002) means that populations in highly modified landscapes are frequently confined to relatively small and geographically distant habitat remnants that experience little if any demographic and genetic rescue because of limited inter-patch immigration (Brown and Kodric-Brown 1977; Brook and Buettel 2016).

Forest remnants resulting from deforestation resemble oceanic islands, so the extinction and colonization processes unified in the equilibrium theory of island biogeography (MacArthur and Wilson 1967) are thought to explain species responses in deforested terrestrial landscapes (Haddad et al. 2015; Resasco et al. 2017). An important difference between oceanic and terrestrial landscapes is that oceanic islands are separated by a truly inhospitable aquatic barrier, whereas forest remnants are generally surrounded by a matrix of modified vegetation that may not be uniformly inhospitable. Many studies have found that the matrix plays an important role in determining patterns of species occurrence, abundance, and richness in forest remnants (Cook et al. 2002; Prevedello and Vieira 2010; Nowakowski et al. 2013a, b). For forest-dwelling species, low- and high-quality matrix may be differentiated by plant biomass (Marzluff and Ewing 2001; Kupfer et al. 2006; Biswas and Wagner 2012). Here we considered high quality matrix to have similar biomass to forest remnants (e.g., secondary regrowth around primary forest remnants), whereas low quality matrix had low plant

biomass (e.g., clear cuts, savanna, or maintained grassland surrounding remnant forest).

Variation in matrix quality can influence organisms in many ways (Kupfer et al. 2006), but movement dynamics are particularly important because they directly influence the colonization success of dispersing individuals (Holderegger and Wagner 2008; Eycott et al. 2010; Martin and Fahrig 2012) that permits demographic and genetic rescue of small populations. Specifically, high matrix quality increases permeability, the extent to which the matrix facilitates movement of individuals among habitat patches (Collinge and Palmer 2002; Haynes and Cronin 2006). For example, ant richness in Mexican coffee plantations (Perfecto and Vandermeer 2002) and squirrel occupancy in urban woodlands in Brussels (Verbeylen et al. 2003) both increased when patches were surrounded by high quality matrix, regardless of the distance to the nearest patch. Populations of Hazel Grouse on forest remnants in Sweden showed evidence of isolation over much shorter distances when surrounded by low-quality matrix (farmland) than high-quality matrix (coniferous forest; Åberg et al. 1995). Arboreal primates in Kenya were encountered more frequently in high-quality matrix (tall vegetation with some tree cover), than in low-quality matrix (short vegetation with little canopy cover; Anderson et al. 2007). The importance of matrix quality is also apparent in synthetic studies across taxa and landscape types. For example, one meta-analysis found that although patch area typically was the primary driver of species occupancy in fragmented habitats, isolation (distance to nearest patch) was most important when habitat patches were embedded in a low-quality, clear-cut matrix (Prugh et al. 2008). Another synthesis found that metrics describing matrix quality were more important for predicting species occupancy and abundance on habitat patches than distance-based isolation metrics (Watling et al. 2011).

Matrix quality has even been shown to influence the strength of the SAR via its relationship with z , the slope of the regression line resulting from $\log_{10}$ transformations of species richness and area of habitat remnants. Generally, z -scores range between 0.1 and 0.5 (Lomolino 2000), with a typical, ‘canonical’ value of 0.262 (Preston 1962). Mean z -scores are greater on oceanic islands than on habitat remnants in agricultural landscapes (Watling and Donnelly 2006), and

greater on forest remnants than in equivalent areas sampled in continuous forest (reviewed in Fahrig 2013). Low z -scores suggest that populations are experiencing low extirpation rates (Losos and Schluter 2000; Rivard et al. 2000; Gao and Perry 2016) at least in part because of demographic and genetic rescue as individuals move through relatively permeable matrix (Hovestadt and Poethke 2005; Kierepka et al. 2016).

Although matrix quality may be a key variable influencing species distributions in modified landscapes, its importance may be taxon-specific, and influenced by additional factors such as the extent of forest cover in a landscape. Species traits may mediate responses to matrix quality because gap-crossing ability (Lees and Peres 2009) or capacity for orienting through the matrix (Pettit et al. 2017) may influence how species perceive matrix quality. With respect to forest cover, the fragmentation threshold hypothesis suggests that negative effects of decreasing patch size and increasing isolation on species richness do not become apparent until 70–90% of habitat has been removed from a landscape (Andrén 1996). When forest cover is high, populations may not be functionally isolated from one another, and species richness may vary little among remnants (Rybicki and Hanski 2013). As forest cover decreases, populations become increasingly isolated, and small populations become vulnerable to extirpation.

It is clear that matrix quality can influence species richness in remnant forest, but the generality and magnitude of matrix effects are not well understood. In an effort to clarify the role of matrix quality in deforested landscapes, we used a meta-analytic approach to address the following research question: How does variation in matrix quality influence the strength of SARs in forest remnants? We predicted that the strength of SARs, measured using an effect size metric derived from the correlation coefficient between species richness and remnant area, would be strongest when matrix quality and forest cover were low. We also predicted the existence of strong SARs in landscapes with low matrix quality for terrestrial dispersers (non-flying invertebrates, amphibians, reptiles, and mammals) that experienced movement constraints and infrequent rescue when matrix quality was low, but weak effects for volant species. Finally, we predicted that matrix quality would be inversely related to the proportion of ice, water, and urban areas in study landscapes, because those land cover types are

thought to be particularly impermeable to movement for many terrestrial species (Forman 2000; Lees and Gilroy 2014). To test our predictions, we coupled SARs from previously published studies from around the world with a metric of matrix quality obtained from remotely sensed satellite data, yielding a synthetic, quantitative assessment of the influence of matrix quality on SARs in remnant forest.

Methods

Literature survey

We searched the primary literature for studies on species richness on forest remnants in modified terrestrial landscapes. To be included in our analysis, we required that studies (1) reported either the correlation coefficient (Pearson's r) between $\log_{10}$ -transformed species richness and remnant area, or raw data from which we could calculate r ; (2) included sufficient information for identification of forest remnants in satellite imagery, either by description, geographic coordinates, or a map of the study area; (3) surveyed forest remnants that could be differentiated from the surrounding landscape in satellite images. If any of the three criteria were not met, the study was excluded from our analysis (Table S1). Two studies investigated different clusters of forest remnants in a single region. The two clusters did not overlap, so were included as separate landscapes, with individual SARs and landscape variables calculated in each case. We focused on remnant forest because of difficulty differentiating non-forest habitat from modified cover in satellite images, and because most studies of habitat modification center on forest.

We first reviewed papers from a previous meta-analysis (Watling and Donnelly 2006), which included 148 studies published through April 2005. Although all of those studies included data on the SAR, only 24 ($\sim 16\%$) met inclusion criteria two and three and were added to our database. We then searched Web of Science with the search terms '*species richness* and *habitat fragmentation* and *isolation*' to update the database with studies from April 2005 to June 2016. Searching these keywords resulted in a total of 339 articles, of which 21 ($\sim 6\%$) met the inclusion criteria. Therefore, our analyses are based on 45 studies conducted in modified landscapes around the

world (Figure S1, Table S2). We extracted the following data from each study: the r value of the correlation between $\log_{10}$ -transformed species richness and remnant area, number of remnants, and taxonomic group (birds, reptiles and amphibians, mammals, plants, or invertebrates).

Describing matrix quality

We downloaded satellite images of each study landscape from the Landsat 4–5 mission archive, which includes data obtained between 1984 and 2013, and used the cloud-free image closest to the date in which fieldwork was completed. All satellite images were obtained within five years of the sampling period. We used standard calibration coefficients (Chandler et al. 2009) to convert the raw data to reflectance values in each of six spectral bands, and projected images into the equal-area Eckert IV projection. We created a minimum bounding polygon circumscribing each study landscape by connecting points superimposed on the satellite images at the outer edges of the most outlying remnants surveyed. To ensure that matrix quality estimates were obtained within a biologically relevant landscape in each study, we buffered each study area polygon by the mean maximum dispersal distance for the study taxon using previously-published data on body mass-dispersal relationships (Jenkins et al. 2007). Buffer radii for each taxonomic group were 32 m for vascular plants, 794 m for inverts, 3.1 km for small rodents, 3.1 km for reptiles and amphibians, 6.3 km for birds, and 63 km for large mammals.

Before assessing matrix quality, it was necessary to differentiate forest remnants from the surrounding matrix. To do this, we created a supervised classification of each study landscape (Churches et al. 2014), using random forest (Breiman 2001) and generalized boosting (Friedman 2001) algorithms to differentiate forest from the surrounding matrix. To create our classification, we first superimposed 1000 points at random throughout each landscape, and visually assessed whether the points intersected forest or non-forest using true- and false-color renderings of the Landsat images. From this pool of classified points, we extracted reflectance values in each of the six spectral bands for 100 randomly selected points in forest and non-forest, respectively. The 200 points were then split randomly into two groups of 100 points each for

model training and testing. A random 75 points from the training set were used to calibrate a model, which was then evaluated using 25 points from the testing set. We repeated this process five times, each time using a unique, random 75–25 training–testing split of the data. Models were evaluated using the true skill statistic (TSS), a metric of model accuracy that ranges from 0 to 1, with higher values indicating greater ability to discriminate forest from non-forest (Allouche et al. 2006). All models has TSS values ≥ 0.70 (Table S2), indicating good classification ability. We therefore used all 200 points to create a prediction map for each landscape. We converted continuous prediction maps indicating the probability that individual cells were forested to binary forest/non-forest maps using a unique threshold for each of the 45 landscapes, determined as the value at which TSS was maximized in the test data.

We used the Enhanced Vegetation Index (EVI) as the basis for our matrix quality metric. The EVI yields values between -1 and 1 , with large values representing high plant biomass (Huete and Justice 1999). The EVI provides accurate estimates of plant biomass even in high-biomass landscapes, and is less prone to atmospheric interference than other indices such as the normalized difference vegetation index (Huete et al. 2002). We calculated a standardized matrix quality metric as

$$1 - \left(\frac{\overline{EVI}_{forest} - \overline{EVI}_{non-forest}}{\overline{EVI}_{total}} \right)$$

by differentiating mean EVI from pixels intersecting forest or non-forest portions of each landscape. The metric describes the difference in plant biomass between forest and matrix as a proportion of the mean EVI in each landscape. Small values of the metric indicate low-quality matrix, where plant biomass was much lower than in forest (Fig S2).

Data analysis

To measure the strength of the SAR in each of the 45 studies, we converted Pearson's r to an estimate of effect size using Fisher's z transformation: $z = 0.5 \times \ln\left(\frac{1+r}{1-r}\right)$ with variance $v_z = \frac{1}{n-3}$. Our first two predictions were evaluated using weighted means analysis of variance (ANOVA), with each study weighted by $\frac{1}{v_z}$. The sum of squares value for each predictor from

weighted means ANOVAs were reported as Cochran's Q heterogeneity statistic, and tested against a Chi square distribution with 1 degree of freedom (Konstantopoulos and Hedges 2009).

We first tested the prediction that the strength of the SAR decreased as matrix quality increased in landscapes with little forest cover. For this test we included matrix quality, forest cover, and their interaction as predictor variables, and Fisher's z as the response variable. We expected this analysis to reveal a significant interaction between matrix quality and forest cover, with particularly strong and negative relationships between effect size and matrix quality in landscapes with low forest cover. Before conducting regressions, we confirmed that matrix quality and forest cover were not highly correlated (Pearson's $r = 0.39$). To test the prediction that the strength of the SAR decreased as matrix quality increased for terrestrial dispersers but not volant taxa, we first categorized the dispersal mode for species in each study as primarily volant (birds and flying invertebrates) or non-volant (all other taxa). We included dispersal mode, matrix quality, and their interaction as predictor variables, and Fisher's z as the response, with the expectation of a significant interaction between matrix quality and dispersal mode. Prior to analysis, we confirmed that the observed number of studies of volant ($N = 19$) and non-volant species ($N = 26$) did not differ significantly from expected counts ($\chi^2 = 1.09$, $df = 1$, $P = 0.297$). Finally, we used simple linear regression to test the prediction that matrix quality decreased with increasing cover of ice, water, and urban areas. Land cover data were acquired from the European Space Agency Climate Change Initiative (ESA 2017) for years 2000, 2005, and 2010, and the layer that corresponded most closely to the year that fieldwork was conducted for each study was used for analysis. Satellite images were processed for analysis using ArcMap version 10.3 (Environmental Systems Research Institute 2012) and all other analyses were completed in R (R core team 2016).

Results

We acquired species and landscape data for 45 landscapes in 24 countries, including 695 patches ranging in size from 0.1 to 23,300 hectares. The 45 studies included in our database were predominately

of invertebrates (16 studies, 36%) and birds (13 studies, 29%), with fewer studies of mammals (nine studies, 20%), amphibians and reptiles (four studies, 9%), and plants (three studies, 7%).

Meta-analysis revealed that the effect of patch area on species richness was influenced by the interaction between matrix quality and forest cover ($Q = 10.75$, $P = 0.001$). Inspection of the interaction plot confirmed that the strength of SARs decreased with increasing matrix quality in low forest cover landscapes, with weaker effects in landscapes with relatively high forest cover (Fig. 1). However, our prediction that matrix quality would interact with dispersal mode to influence the strength of SARs was not supported ($Q = 1.01$, $P = 0.315$). There was a significant main effect of dispersal mode on the strength of SARs ($Q = 18.26$, $P < 0.001$), with overall stronger SARs reported for volant compared with non-volant species (Fig. 2). As expected from our third prediction, matrix quality decreased as landscapes became increasingly dominated by impermeable land cover types (water, ice, and urban areas; $F_{1,43} = 17.94$, $P < 0.001$; Fig. 3).

Discussion

Here we provide clear, quantitative evidence to suggest that matrix quality plays a major role in determining patterns of species richness in remnant forest. Species-area relationships were particularly strong in landscapes with low matrix quality and little remaining forest. We found that SARs were stronger for volant species such as birds and flying invertebrates than for terrestrial dispersers, but no differences in the response to matrix quality by dispersal mode. Our data underscore the importance of conservation actions that focus on increasing matrix quality, such as actively or passively promoting vegetation regrowth in deforested landscapes.

We found that the strength of the SAR decreased with increasing matrix quality in landscapes with less than about 20% forest cover (Fig. 1). The relationship implies that low matrix quality may exacerbate the negative effects of habitat loss in severely deforested landscapes. The matrix matters for biodiversity in part because it influences the rate and success of movement among habitat remnants (Prugh et al. 2008; Prevedello and Vieira 2010; Ruffell et al. 2017). For example,




Fig. 1 The strength of the species-area relationship (measured as the effect size of the correlation between $\log_{10}$ (patch area) and $\log_{10}$ (species richness)) decreased with matrix quality in landscapes with low forest cover, but was less affected by matrix quality in landscapes with high forest cover. Here we describe matrix quality in terms of plant biomass, such that high-quality matrix has greater plant biomass than low-quality

matrix. To illustrate the interaction between forest cover and matrix quality on the effect size of the species-area relationship, we modeled the relationship between Fisher's z and matrix quality at three levels of forest cover: 10, 20, and 30%. Only four of 45 studies included in the analysis occurred in landscapes where forest cover exceeded 30%

voles in Indiana meadows (Russell et al. 2007) and beetles in California prairies (Collinge and Palmer 2002) have been found to avoid or make reduced use of portions of the matrix comprised of low-quality, short-statured vegetation compared with tall grasses that more closely resemble their forest habitat. In another study, understory forest birds avoided corn fields, preferring to move among forest remnants via relatively high-quality *Eucalyptus* stands, possibly because of decreased predation risk or increased resource availability (Biz et al. 2017). When large differences in plant biomass between forest and matrix in low-quality landscapes prevent inter-patch movement and decrease functional connectivity (Tischendorf and Fahrig 2000; Smith and Hellmann 2002; Vasudev et al. 2015), remnants may experience infrequent demographic or genetic rescue (Brown and Kodric-Brown 1977), a process that is likely to be

particularly important in the most severely deforested landscapes (Fig. 1).

We observed stronger overall SARs for volant compared with non-volant taxa, but no indication that responses to matrix quality differed significantly by dispersal mode. Although we were initially surprised that SARs were strongest for volant species, previous research has demonstrated that both birds and flying invertebrates may limit travel through the matrix (Desrochers and Hannon 1997; Castellón and Sieving 2006). Some species of understory birds even avoid entirely crossing the matrix (Harris and Reed 2002; Şekercioğlu et al. 2002). Furthermore, birds and butterflies often respond relatively quickly to even subtle structural changes in the matrix, suggesting that they have great ability to perceive matrix quality (Ricketts 2001; Ries and Debinski 2001; Martin and Possingham 2005). Inter-patch movements may be

Fig. 2 The mean effect of patch area on species richness was greater for volant species than for non-volant taxa. Individual effects for each study are superimposed over the boxplots





Fig. 3 Matrix quality decreases as the proportion of impermeable terrestrial cover types (ice, water, and urban areas) increases



severely or entirely inhibited for species able to sample and perceive low-quality matrix as inhospitable (Clobert et al. 2009), whereas species with limited perceptual ability may not avoid the matrix as frequently. Although decades of research have revealed few taxon-based differences in SARs that can be interpreted clearly in terms of dispersal ability (Öckinger et al. 2010; Aranda et al. 2013; Matthews et al. 2016; Fattorini et al. 2017), most of that work has focused on *z*-scores, rather than the effect size metric we analyzed here. We suggest that the strong effect of patch size on species richness that we observed for volant species may be a consequence of greater perceptual abilities for flying organisms compared with passive or terrestrial dispersers, but emphasize that more data are needed to evaluate this possibility.

Our analysis found particularly strong gradients in species richness in landscapes where forest remnants were surrounded by low-quality matrix. The most important way to mitigate species losses in modified landscapes is to minimize deforestation, although alleviating the pace of deforestation is complex, especially in the face of a growing human population and the possibility of future food insecurity (Garibaldi et al. 2017). However, our results imply that minimizing species losses in deforested landscapes may also be achieved by enhancing matrix quality to increase landscape connectivity. Several features have been suggested to increase landscape connectivity, including corridors (Gilbert-Norton et al. 2010; Haddad et al. 2017) and small stepping stone fragments that reduce effective isolation in fragmented landscapes (Baum et al. 2004; Saura 2014). Another way to increasing landscape connectivity is to increase matrix permeability via passive or active regeneration (Smallbone et al. 2014; de Rezende et al. 2015). Although active regeneration of the matrix may be more expensive than passive regeneration (Brancalion et al. 2016), it may increase the value of ecosystem services in managed landscapes, partially compensating for production losses tied to livestock, agriculture, and timber harvest (Bullock et al. 2011).

Managing the matrix to avert species losses in modified landscapes has become an increasingly important conservation strategy. One of the best examples of a large-scale conservation strategy focused on enhancing matrix quality is the Atlantic Forest Restoration Pact, in which more than 260 groups have organized to actively restore 15 million

hectares of deforested and degraded lands by the year 2050 with the goal of conserving biodiversity (Pinto et al. 2014). In Costa Rica, the Payment for Environmental Services Program reimburses landowners for the ecosystem services their property provides through biodiversity, water, and carbon payments, as long as landowners protect existing vegetation or plant tree plantations on their property (Pagiola 2008). The benefit of conservation plans such as these are multi-faceted, enhancing the delivery of ecosystem services while increase landscape connectivity, with collateral benefits for biodiversity by facilitating movement between forest remnants.

We used satellite imagery and remote sensing techniques to provide a metric of matrix quality that helps explain trends in species responses to landscape modification. A drawback to using a single metric is that it may not adequately describe matrix quality for all species. For example, distributions of some species are limited by the presence of keystone species rather than habitat quality (e.g., presence of predators determined by prey; Delibes-Mateos et al. 2007). Also, resource-limited habitat specialists such as monarch butterflies may not use the matrix the same way as generalist butterflies (Flockhart et al. 2017). Our approach does not provide a substitute for detailed species management plans. However, because it is impractical to generate species-specific data for all species of conservation concern, our approach to assessing matrix quality may be applied as an initial assessment for many species at once. Data generalizing species responses to matrix quality can be used to suggest preliminary conservation strategies for many, rather than few species where immediate conservation practices are needed (Lambeck 1997). For example, in Peninsular Malaysia, 558 isolated karst forests are at risk of destruction, and in just twelve forests, sixteen endemic and seven karst forest-adapted reptile species have been discovered in a seven-year period (Grismer et al. 2016). It is likely that these Malaysian species, as well as many others, are at risk of extinction without immediate conservation action. Our data suggest that efforts to increase matrix quality should reduce the threat of extirpation of vulnerable populations on small forest remnants.

In conclusion, we found that the effect of patch area on species richness is greatest in severely deforested landscapes where matrix quality is low (i.e., there is a large difference in plant biomass between forest

remnants and the surrounding matrix). Our analysis of global land cover confirmed that matrix quality was lowest in areas dominated by three relatively impermeable cover types: ice, water, and urban land. The impact of matrix quality on species-area relationships was particularly strong for volant species. Although the remotely sensed matrix quality metric we used here is not a perfect substitute for field data describing how plant biomass influences organism movement and population persistence, it does provide an approach that can be used to generalize one type of matrix effect on populations in remnant forest. A key implication of our work is that conservation strategies that incentivize active or passive restoration of the matrix should help maintain populations on forest remnants.

Acknowledgements We thank R Drenovsky, C Anthony, and D Hickman for constructive feedback on the manuscript, and the Department of Biology at John Carroll University for supporting IJR during data collection and writing of this paper.

References

- Åberg J, Jansson G, Swenson JE, Angelstam P (1995) The effect of matrix on the occurrence of hazel grouse (*Bonasa bonasia*) in isolated habitat fragments. *Oecologia* 103:265–269
- Allouche O, Tsoar A, Kadmon R (2006) Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *J Appl Ecol* 43:1223–1232
- Anderson J, Rowcliffe JM, Cowlishaw G (2007) Does the matrix matter? A forest primate in a complex agricultural landscape. *Biol Conserv* 135:212–222
- Andrén H (1996) Population responses to habitat fragmentation: statistical power and the random sample hypothesis. *Oikos* 76:235–242
- Aranda SC, Gabriel R, Borges PAV, Santos AMC, Hortal J, Baselga A, Lobo JM (2013) How do different dispersal models shape the species-area relationship? Evidence for between-group coherence in the Macaronesian flora. *Glob Ecol Biogeogr* 22:483–493
- Barnosky AD, Hadly EA, Bascompte J, Berlow EL, Brown JH, Fortelius M, Getz WM, Harte J, Hastings A, Marquet PA, Martinez ND, Mooers A, Roopnarine P, Vermeij G, Williams JW, Gillespie R, Kitzes J, Marshall C, Matzke N, Mindell DP, Revilla E, Smith AB (2012) Approaching a state shift in Earth's biosphere. *Nature* 486:52–58
- Baum KA, Haynes KJ, Dilleuth FP, Cronin JT (2004) The matrix enhances the effectiveness of corridors and stepping stones. *Ecology* 85:2671–2676
- Biswas SR, Wagner HH (2012) Landscape contrast: a solution to hidden assumptions in the metacommunity concept? *Landscape Ecol* 27:621–631
- Biz M, Cornelius C, Paul J, Metzger W (2017) Matrix type affects movement behavior of a Neotropical understory forest bird. *Perspect Ecol Conserv* 15:10–17
- Brançalion PHS, Schweizer D, Gaudare U, Mangueira JR, Lamonato F, Farah FT, Nave AG, Rodrigues RR (2016) Balancing economic costs and ecological outcomes of passive and active restoration in agricultural landscapes: the case of Brazil. *Biotropica* 48:856–867
- Breiman L (2001) Random forests. *Mach Learn* 45:5–32
- Brook BW, Buettel JC (2016) Emigration is costly, but immigration has benefits in human-altered landscapes. *Funct Ecol* 30:1478–1479
- Brown JH, Kodric-Brown A (1977) Turnover rates in insular biogeography: effect of immigration on immigration on extinction. *Ecology* 58:445–449
- Bullock JM, Aronson J, Newton AC, Pywell RF, Rey-benayas JM (2011) Restoration of ecosystem services and biodiversity: conflicts and opportunities. *Trends Ecol Evol* 26:541–549
- Castellón TD, Sieving KE (2006) An experimental test of matrix permeability and corridor use by an endemic understory bird. *Conserv Biol* 20:135–145
- Chandler G, Markham BL, Helder DL (2009) Summary of current radiometric calibration coefficients for Landsat MSS, TM, ETM+, and EO-1 ALI sensors. *Remote Sens Environ*
- Churches CE, Wampler PJ, Sun W, Smith AJ (2014) Evaluation of forest cover estimates for Haiti using supervised classification of Landsat data. *Int J Appl Earth Obs Geoinf* 30:203–216
- Clobert J, Le Galliard JF, Cote J, Meylan S, Massot M (2009) Informed dispersal, heterogeneity in animal dispersal syndromes and the dynamics of spatially structured populations. *Ecol Lett* 12:197–209
- Collinge SK, Palmer TM (2002) The influences of patch shape and boundary contrast on insect response to fragmentation in California grasslands. *Landscape Ecol* 17:647–656
- Cook WM, Lane KT, Foster BL, Holt RD (2002) Island theory, matrix effects and species richness patterns in habitat fragments. *Ecol Lett* 5:619–623
- de Rezende CL, Uezu A, Scarano FR, Araujo DSD (2015) Atlantic Forest spontaneous regeneration at landscape scale. *Biodivers Conserv* 24:2255–2272
- Delibes-Mateos M, Redpath SM, Angulo E, Ferreras P, Villafuerte R (2007) Rabbits as a keystone species in southern Europe. *Biol Conserv* 137:149–156
- Desrochers A, Hannon SJ (1997) Gap crossing decisions by forest songbirds during the post-fledging period. *Conserv Biol* 11:1204–1210
- Edgar GJ, Stuart-Smith RD, Willis T, Kininmonth S, Baker SC, Banks S, Barrett NS, Becerro MA, Bernard ATF, Berkhout J, Buxton CD, Campbell SJ, Cooper AT, Davey M, Edgar SC, Försterra G, Galván DE, Irigoyen AJ, Kushner DJ, Moura R, Ed Parnell P, Shears NT, Soler G, Strain SMA, Thomson RJ (2014) Global conservation outcomes depend on marine protected areas with five key features. *Nature* 506:216–220
- Environmental Systems Research Institute (2012) ArcGIS Release 10.3. Redlands, California
- ESA (2017) Land Cover CCI Product User Guide Version 2.0. <http://www.fao.org/3/a-i4808e.pdf>. Accessed 14 Feb 2017

- Eycott A, Watts K, Brandt G, Buyung-Ali L, Bowler D, Stewart G, Pullin AS (2010) Do landscape matrix features affect species movement? Collaboration for Environmental Evidence review 08-006 (SR 43)
- Fahrig L (2003) Effects of habitat fragmentation on biodiversity. *Annu Rev Ecol Evol Syst* 34:487–515
- Fahrig L (2013) Rethinking patch size and isolation effects: the habitat amount hypothesis. *J Biogeogr* 40:1649–1663
- Fattorini S, Borges PAV, Dapporto L, Strona G (2017) What can the parameters of the species-area relationship (SAR) tell us? Insights from Mediterranean islands. *J Biogeogr* 44:1018–1028
- Flockhart DTT, Brower LP, Ramirez MI, Hobson KA, Wassenaar LI, Altizer S, Norris DR (2017) Regional climate on the breeding grounds predicts variation in the natal origin of monarch butterflies overwintering in Mexico over 38 years. *Glob Change Biol* 23:2565–2576
- Forman RTT (2000) Estimate of the area affected ecologically by the road system in the United States. *Conserv Biol* 14:31–35
- Friedman JH (2001) Greedy function approximation: a gradient boosting machine. *Ann Stat* 29:1189–1232
- Gao D, Perry G (2016) Species-area relationships and additive partitioning of diversity of native and nonnative herpetofauna of the West Indies. *Ecol Evol* 6:7742–7762
- Garibaldi LA, Gemmill-Herren B, D'Annolfo R, Graeb BE, Cunningham SA, Breeze TD (2017) Farming approaches for greater biodiversity, livelihoods, and food security. *Trends Ecol Evol* 32:68–80
- Gilbert-Norton L, Wilson R, Stevens JR, Beard KH (2010) A meta-analytic review of corridor effectiveness. *Conserv Biol* 24:660–668
- Gonzalez A, Chaneton EJ (2002) Heterotroph species extinction, abundance and biomass dynamics in an experimentally fragmented microecosystem. *J Anim Ecol* 71:594–602
- Grismer JL, Schulte JA II, Alexander A, Wagner P, Travers SL, Buehler MD, Welton LJ, Brown RM (2016) The Eurasian invasion: phylogenomic data reveal multiple Southeast Asian origins for Indian Dragon Lizards. *BMC Evol Biol* 16:43
- Haddad NM, Brudvig LA, Clobert J, Davies KF, Conzaes A, Holt RD, Lovejoy TE, Sexton JO, Austin MP, Collins CD, Cook WM, Damschen EI, Ewers RM, Foster BL, Jenkins CN, King AJ, Laurance WF, Levey DJ, Margules CR, Melbourne BA, Nicholls AO, Orrock JL, Song d-X, Townsend JR (2015) Habitat fragmentation and its lasting impact on Earth's ecosystems. *Sci Adv* 1:e1500052
- Haddad NM, Gonzalez A, Brudvig LA, Burt MA, Levey DJ, Damschen EI (2017) Experimental evidence does not support the habitat amount hypothesis. *Ecography* 40:48–55
- Haila Y (2002) A conceptual genealogy of fragmentation research: from island biogeography to landscape ecology. *Ecol Appl* 12:321–334
- Harris RJ, Reed JM (2002) Behavioral barriers to non-migratory movements of birds. *Ann Zool Fenn* 39:275–290
- Haynes KJ, Cronin JT (2006) Interpatch movement and edge effects: the role of behavioral responses to the landscape matrix. *Oikos* 113:43–54
- Holderegger R, Wagner HH (2008) *Landsc Genet*. Bioscience 58:199–207
- Hovestadt T, Poethke HJ (2005) Dispersal and establishment: spatial patterns and species-area relationships. *Divers Distrib* 11:333–340
- Huete A, Didan K, Miura T, Rodriguez EP, Gao X, Ferreira LG (2002) Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sens Environ* 83:195–213
- Huete AR, Justice C (1999) MODIS vegetation index (MOD13) algorithm theoretical basis document
- Jenkins DG, Brescacin CR, Duxbury CV, Elliott JA, Evans JA, Grablow KR, Hillegass M, Lyon BN, Metzger GA, Olandese ML, Pepe D, Silvers GA, Suresch HN, Thompson TN, Trexler CM, Williams GE, Williams NC, Williams SE (2007) Does size matter for dispersal distance? *Glob Ecol Biogeogr* 16:415–425
- Jones NT, Germain RM, Grainger TN, Hall AM, Baldwin L, Gilbert B (2015) Dispersal mode mediates the effect of patch size and patch connectivity on metacommunity diversity. *J Ecol* 103:935–994
- Kierepka E, Anderson SJ, Swihart RK, Rhodes OE Jr (2016) Evaluating the influence of life-history characteristics on genetic structure: a comparison of small mammals inhabiting complex agricultural landscapes. *Ecol Evol* 6:6376–6396
- Konstantopoulos S, Hedges LV (2009) Analyzing effect sizes: fixed-effects models. In: Cooper H, Hedges LV, Valentine JC (eds) *The handbook of research synthesis and meta-analysis*. Russell Sage Foundation, New York, pp 279–293
- Kupfer JA, Malanson GP, Franklin SB (2006) Not seeing the ocean for the islands: the mediating influence of matrix-based processes on forest fragmentation effects. *Glob Ecol Biogeogr* 15:8–20
- Lambeck RJ (1997) Focal species: a multi-species umbrella for nature conservation. *Conserv Biol* 11:849–856
- Lees AC, Gilroy JJ (2014) Vagrancy fails to predict colonization of oceanic islands. *Glob Ecol Biogeogr* 23:405–441
- Lees AC, Peres CA (2009) Gap-crossing movements predict species occupancy in Amazonian forest fragments. *Oikos* 118:280–290
- Lomolino MV (2000) Ecology's most general, yet protean pattern: the species-area relationship. *J Biogeogr* 27:17–26
- Losos BJ, Schluter D (2000) Analysis of an evolutionary species-area relationship. *Nature* 408:847–850
- MacArthur RH, Wilson EO (1967) *The Theory of Island Biogeography*. Princeton University Press, New Jersey
- Martin AE, Fahrig L (2012) Measuring and selecting scales of effect for landscape predictors in species-habitat models. *Ecol Appl* 22:2277–2292
- Martin TG, Possingham HP (2005) Predicting the impact of livestock grazing on birds using foraging height data. *J Appl Ecol* 42:400–408
- Marzluff JM, Ewing K (2001) Restoration of fragmented landscapes for the conservation of birds: a general framework and specific recommendations for urbanizing landscapes. *Restor Ecol* 9:280–292
- Matthews TJ, Guilhaumon F, Triantis KA, Borregaard MK, Whittaker RJ (2016) On the form of species-area relationships in habitat islands and true islands. *Glob Ecol Biogeogr* 25:847–858

- Newbold T, Hudson LN, Arnell AP, Contu S, De Palma A, Ferrier S, Hill SLL, Hoskins AJ, Lysenko I, Phillips HRP, Burton VJ, Chng CWT, Emerson S, Gao D, Pask-Hale G, Hutton J, Jung M, Sanchez-Ortiz K, Simmons BI, Whitmee S, Zhang H, Scharlemann JPW, Purvis A (2016) Has land use pushed terrestrial biodiversity beyond the planetary boundary? A global assessment. *Science* 353:288–291
- Nowakowski AJ, Hyslop N, Watling JJ, Donnelly MA (2013a) Matrix type alters structure of aquatic vertebrate assemblages in cypress domes. *Biodivers Conserv* 22:497–511
- Nowakowski AJ, Otero Jiménez B, Allen M, Diaz-Escobar M, Donnelly MA (2013b) Landscape resistance to movement of the poison frog, *Oophaga pumilio*, in the lowlands of northeastern Costa Rica. *Anim Conserv* 16:188–197
- Öckinger E, Schweiger O, Crist TO, Debinski DM, Krauss J, Kuussaari M, Peterson JD, Pöyry J, Settele J, Summerville KS, Bommarco R (2010) Life-history traits predict species responses to habitat area and isolation: a cross-continental synthesis. *Ecol Lett* 13:969–979
- Pagiola S (2008) Payments for environmental services in Costa Rica. *Ecol Econ* 65:712–724
- Paz Durán A, Inger R, Cantú-Salazar L, Gaston KJ (2016) Species richness representation with protected areas is associated with multiple interacting spatial features. *Divers Distrib* 22:300–308
- Perfecto I, Vandermeer J (2002) Quality of agroecological matrix in a tropical montane landscape: ants in coffee plantations in Southern Mexico. *Conserv Biol* 16:174–182
- Pettit L, Greenlees M, Shine R (2017) The impact of transport and translocation on dispersal behaviour in the invasive cane toad. *Oecologia* 184:411–422
- Pinto SR, Melo F, Tabarelli M, Padovesi A, Carrascosa H, Calmon M, Rodrigues R, Gomes César R, Brancalion PHS (2014) Governing and delivering a biome-wide restoration initiative: the case of Atlantic Forest Restoration Pact in Brazil. *Forests* 5:2212–2229
- Preston FW (1962) The canonical distribution of commonness and rarity: part I. *Ecology* 43:187–215
- Prevedello JA, Vieira MV (2010) Does the type of matrix matter? A quantitative review of the evidence. *Biodivers Conserv* 19:1205–1223
- Prugh LR, Hodges KE, Sinclair ARE, Brashares JS (2008) Effect of habitat area and isolation on fragmented animal populations. *Proc Natl Acad Sci USA* 105:20770–20775
- R Core Team (2016) R: A language and environment for statistical computing. R foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>
- Resasco J, Bruna EM, Haddad NM, Banks-Leite C, Margules CR (2017) The contribution of theory and experiments to conservation in fragmented landscapes. *Ecography* 40:109–118
- Ricketts TH (2001) The matrix matters: effective isolation in fragmented landscapes. *Am Nat* 158:87–99
- Ries L, Debinski DM (2001) Butterfly responses to habitat edges in the highly fragmented prairies of central Iowa. *J Anim Ecol* 70:840–852
- Rivard DH, Poitevin J, Plasse D, Carleton M, Currie DJ (2000) Changing species richness and composition in Canadian national parks. *Conserv Biol* 14:1099–1109
- Ruffell J, Clout MN, Didham RK (2017) The matrix matters, but how should we manage it? Estimating the amount of high-quality matrix required to maintain biodiversity in fragmented landscapes. *Ecography* 40:171–178
- Russell RE, Swihart RK, Craig BA (2007) The effects of matrix structure on movement decisions of meadow voles (*Microtus pennsylvanicus*). *J Mammal* 88:573–579
- Rybicki J, Hanski I (2013) Species-area relationships and extinctions caused by habitat loss and fragmentation. *Ecol Lett* 16:27–38
- Saura S (2014) Stepping stones are crucial for species' long-distance dispersal and range expansion through habitat networks. *Appl Ecol* 51:171–182
- Şekercioğlu CH, Ehrlich PR, Daily GC, Goehring D, Sandi R (2002) Disappearance of insectivorous birds from tropical forest fragments. *Proc Natl Acad Sci USA* 99:263–267
- Smallbone LT, Matthews A, Lunt ID (2014) Landscape and Urban Planning regrowth provides complementary habitat for woodland birds of conservation concern in a regenerating agricultural landscape. *Landsc Urban Plan* 124:43–52
- Smith JNM, Hellmann JJ (2002) Population persistence in fragmented landscapes. *Trends Ecol Evol* 17:397–399
- Steinmann K, Eggenberg S, Wohlgemuth T, Linder HP, Zimmermann NE (2011) Niches and noise-disentangling habitat diversity and area effect on species diversity. *Ecol Complex* 8:313–319
- Tischendorf L, Fahrig L (2000) On the usage and measurement of landscape connectivity. *Oikos* 90:7–19
- Vasudev D Jr, Fletcher Jr RJ, Goswami VR, Krishnadas M (2015) From dispersal constraints to landscape connectivity: lessons from species distribution modeling. *Ecography* 38:967–978
- Verbeyleen G, De Bruyn L, Adriaensen F, Matthysen E (2003) Does matrix resistance influence Red squirrel (*Sciurus vulgaris* L. 1758) distribution in an urban landscape? *Landscape Ecol* 18:791–805
- Watling JJ, Donnelly MA (2006) Fragments as islands: a synthesis of faunal responses to habitat patchiness. *Conserv Biol* 20:1016–1025
- Watling JJ, Nowakowski AJ, Donnelly MA, Orrock JL (2011) Meta-analysis reveals the importance of matrix composition for animals in fragmented habitat. *Glob Ecol Biogeogr* 20:209–217
- Watson JEM, Dudley N, Segan DB, Hockings M (2014) The performance and potential of protected areas. *Nature* 515:67–73