

IN DEPTH

Socioeconomic Status and Cardiovascular Outcomes

Challenges and Interventions

ABSTRACT: Socioeconomic status (SES) has a measurable and significant effect on cardiovascular health. Biological, behavioral, and psychosocial risk factors prevalent in disadvantaged individuals accentuate the link between SES and cardiovascular disease (CVD). Four measures have been consistently associated with CVD in high-income countries: income level, educational attainment, employment status, and neighborhood socioeconomic factors. In addition, disparities based on sex have been shown in several studies. Interventions targeting patients with low SES have predominantly focused on modification of traditional CVD risk factors. Promising approaches are emerging that can be implemented on an individual, community, or population basis to reduce disparities in outcomes. Structured physical activity has demonstrated effectiveness in low-SES populations, and geomapping may be used to identify targets for large-scale programs. Task shifting, the redistribution of healthcare management from physician to nonphysician providers in an effort to improve access to health care, may have a role in select areas. Integration of SES into the traditional CVD risk prediction models may allow improved management of individuals with high risk, but cultural and regional differences in SES make generalized implementation challenging. Future research is required to better understand the underlying mechanisms of CVD risk that affect individuals of low SES and to determine effective interventions for patients with high risk. We review the current state of knowledge on the impact of SES on the incidence, treatment, and outcomes of CVD in high-income societies and suggest future research directions aimed at the elimination of these adverse factors, and the integration of measures of SES into the customization of cardiovascular treatment.

William M. Schultz, MD
Heval M. Kelli, MD
John C. Lisko, MD
Tina Varghese, MD
Jia Shen, MD
Pratik Sandesara, MD
Arshed A. Quyyumi, MD
Herman A. Taylor, MD
Martha Gulati, MD
John G. Harold, MD
Jennifer H. Mieres, MD
Keith C. Ferdinand, MD
George A Mensah, MD
Laurence S. Sperling, MD

Key Words: cardiovascular diseases
■ epidemiology ■ social class ■ social
determinants of health ■ risk factors

© 2018 American Heart Association, Inc.

<http://circ.ahajournals.org>

Cardiovascular diseases (CVDs) remain the leading cause of death from chronic disease in the United States and worldwide despite remarkable advances made in the past century. CVD accounts for nearly 1 in 3 deaths in the United States despite a 25.3% decrease in age-standardized deaths attributed to CVD from 2004 to 2014.¹ Traditional CVD risk factors have been identified, including hypertension, dyslipidemia, diabetes mellitus, family history of premature coronary heart disease (CHD), and smoking; however, the contribution of social determinants of health, as best represented by socioeconomic status (SES), to CVD risk is poorly understood. Low SES has been linked to the development of CVD and may confer a cardiovascular risk that is equivalent to traditional risk factors.^{2,3} The increased burden of CVD in people with low SES is attributable to a constellation of biological, behavioral, and psychosocial risk factors that are more prevalent in disadvantaged individuals.^{4,5}

Established interventions exist for addressing SES-related determinants of health and health inequities.⁶ Several interventions have been evaluated to improve health disparities attributable to social inequity, but results are inconsistent.⁷ Better understanding of SES and risk stratification are critical first steps in identifying and tailoring interventions to improve CVD risk in the population. We review the current state of knowledge on the impact of SES on the incidence, treatment, and outcomes of CVD in high-income societies and suggest future research directions aimed at the elimination of these adverse factors, and the integration of measures of SES into the customization of cardiovascular treatment, as well.

OVERVIEW OF SOCIOECONOMIC FACTORS

The factors that constitute an individual's SES vary between location and culture.^{8–10} Four markers for SES have demonstrated an association with CVD in high-income countries (HIC): income level, educational attainment, employment status, and environmental factors (Figure 1).^{11–17} Although many factors have been used as surrogates for SES, these particular markers are advantageous because of their ease of collection by questionnaire in addition to the substantial existing literature characterizing their association with clinical CVD outcomes. In addition, it is unlikely that a single marker of SES is adequate in predicting CVD risk because of the interactions between SES factors, regional variations in SES associations with CVD, and the dynamic changes between SES markers and CVD outcomes throughout an individual's life course.^{8,11,18}

Low- and middle-income countries (LMICs) carry ≈80% of the global burden of CVD.¹⁹ However, studies




Figure 1. Markers of socioeconomic status.

evaluating the association between SES and cardiovascular health in LMICs are limited and often contain conflicting findings.^{10,20} Furthermore, the association of SES and CVD in LMICs cannot be extrapolated from studies in HICs; obesity is an epidemic among the impoverished in HICs, but is a disease of the rich in low-income countries because of the increased access to a Western diet.²¹ Characterization of the impact of SES in LMICs will require additional high-quality prospective studies and is beyond the scope of this review.

INCOME LEVEL

Income level has been consistently associated with CVD risk.^{11,12} A large study in the United States and Finland found an increased risk of nonfatal myocardial infarction and sudden cardiac death in the low-income cohorts that persisted after adjusting for smoking and alcohol consumption.¹² The results may be applicable at both an individual and neighborhood level.²² Gerber et al²² found that each \$10 000 increase in median income of a neighborhood reduced mortality risk in the group by 10%.

A study of >15 000 patients admitted for acute myocardial infarction (AMI) or CHD in the Netherlands found that individuals in the lower quintiles of income had significantly higher rates of 28-day and 1-year mortality.²³ Increased mortality was also shown for the lowest-income quartile in comparison with the highest-income quartile in a cohort presenting with ST-segment-elevation myocardial infarction (hazard ratio, 1.17; 95% confidence interval, 1.11–1.25).²⁴ Residents in low-income areas were less likely to receive a left heart catheterization within 24 hours of a ST-segment-elevation myocardial infarction or within

48 hours of a non-ST-segment-elevation myocardial infarction than patients from high-income areas.²⁴ The authors of this study posited that the difference may be attributable to the perception that low-income individuals would be less likely to comply with critical post-percutaneous coronary intervention medications (eg, antiplatelet agents) or that low-income patients were less likely to be offered more expensive procedures (eg, drug-eluting stents).²⁴ The mortality difference between income groups in the study was attenuated after adjustment for time to left heart catheterization.²⁴ Increased mortality rates were also present in patients from low-income areas in Taiwan, a HIC with universal health care, who were hospitalized with AMI.²⁵

Mortality differences in low-income patients may be driven in part by disparities in standards of care. Low-income patients in the Netherlands were less likely to undergo interventions, including percutaneous coronary intervention, when presenting with AMI.²³ Cardiac rehabilitation, a proven therapy to reduce mortality and hospital readmissions after AMI,^{26,27} is less likely to be attended or completed by individuals from low-income areas after hospitalization for CHD, percutaneous coronary intervention, or coronary artery bypass graft surgery.²⁸ Studies have also demonstrated that statin medications were less likely to be initiated after AMI in low-income individuals.^{29,30} Men with low incomes were less likely to receive other guideline-recommended medications, including β -blockers and angiotensin-converting enzyme inhibitors, but the association was attenuated in women.³⁰

The substandard care may be explained in part by decreased access to quality care in the socially disadvantaged.^{31–33} Lack of access to quality care has been shown to increase heart failure and post-AMI hospital readmission rates in the United States,³³ and patients with financial barriers to health care had fewer yearly medical checkups, were less likely to use aspirin, and were at higher risk of future CVD events.³²

Low-income individuals in Taiwan undergoing coronary artery bypass graft received lower-quality care and experienced higher mortality than the high-income cohort.³¹ France, like Taiwan, has a national health system but still faces inequalities in access to primary care and coronary revascularizations in low-income areas.³⁴ These studies highlight the difficulty in providing equal care to a patient population regardless of income. Inequalities in access to care, interventions, and prescribing practices will need to be addressed before care becomes equivalent between social groups.

EDUCATIONAL ATTAINMENT

The inverse relationship between educational attainment and CVD in HIC has been known for decades,

and new population-based studies continue to provide insight into the strength of the association and its underlying mechanisms.^{13,35} A large study by Woodward et al³⁵ analyzed nearly 90 000 individuals in Australia and New Zealand and found that those with a primary education had an increased risk of CVD, cardiovascular mortality, and all-cause mortality in comparison with individuals with a tertiary education. Higher education was associated with increased alcohol consumption and inversely related to smoking, blood pressure, cholesterol levels, and diabetes mellitus.³⁵

Several studies have also demonstrated a higher risk of AMI in individuals with a lower educational level.^{36–38} Lower educational attainment also predicts worse short-term (30 days) and long-term (≥ 1 year) outcomes after AMI.^{22,39–41} Of the several socioeconomic factors examined in a cohort of patients from South Korea, only low education (≤ 6 years of schooling) was associated with increased risk of cardiac events or all-cause mortality.⁴¹ Patients with low SES tend to have an increased number of comorbidities but receive fewer interventions.^{40,42} Despite their increased comorbidities, individuals with a lower education level were less likely to be referred to tertiary prevention programs, including cardiac rehabilitation, than those who had a higher education level.⁴²

Educational attainment may affect health in several ways. Individuals with less education tend to have an increased number of CVD risk factors.⁵ An analysis in the Netherlands by Kershaw et al⁴³ demonstrated that a majority (56.6%) of CHD risk in individuals with low education was attributable to behavioral and biological risk factors. The most significant contributors were smoking (27.3%), obesity (10.2%), physical inactivity (6.3%), and hypertension (5.3%).⁴³ Hu et al³⁶ determined that approximately half of the increased risk of incident AMI in low education groups was explained by traditional risk factors. Even with these estimates, the mechanisms underlying the remainder of the increased risk associated with low educational attainment remain to be determined.

A potential contributor to CVD risk is the strong correlation between education and health literacy.^{44,45} Individuals with poor health literacy are more likely to be noncompliant with their medications⁴⁶ and experience increased all-cause mortality.⁴⁷ The association between health literacy and outcomes may be partially mediated by poor reading comprehension, which has an indirect effect on CHD risk.⁴⁸ The absolute effect of health literacy and reading comprehension on CVD risk requires further characterization, but may be an area for intervention in individuals with low education attainment. It is also important to note that although studies looking at the impact of SES on outcomes adjust for variables such as traditional risk factors, there is potential for residual confounding that may be contributing to adverse health that is not accounted for in the models.

EMPLOYMENT STATUS

Employment is a commonly used marker of SES, and unemployment has been associated with increased risk of CVD.^{11,14,49} Analysis by Méjean et al⁴⁹ demonstrated a 20% increase in risk of CHD events in an unemployed French population without preexisting heart disease (hazard ratio, 1.20; confidence interval, 1.04–1.39) after adjustment for age, sex, diet, and lifestyle. A large proportion (46%) of the CHD risk is explained by dietary and lifestyle mediators, most notably alcohol consumption (17%) and smoking (13%).⁴⁹

A recent study of individuals with high SES relative to the general population also demonstrated an increased risk of cardiovascular events in the unemployed despite extensive adjustment for covariates, including age, sex, biological characteristics, behavioral variables, and socioeconomic factors.¹⁴ The outcomes of the unemployed population in the study were worse than those of the retired cohort, which suggests that the detrimental effect of unemployment may be driven by the job loss itself.¹⁴ An alternative explanation is that those with poor health may be more likely to become unemployed. Nonetheless, it is important to note that unemployment is associated with increased CVD burden in the socially advantaged, and in those with low SES at baseline, as well.

Dupre et al⁵⁰ performed a comprehensive study on the relationship between employment status and risk for AMI in the United States. Unemployment was associated with a 35% increased risk for AMI (hazard ratio, 1.35; confidence interval, 1.10–1.66) within the first year, after which time the risk disappeared.⁵⁰ The cumulative number of job losses was associated with an incremental increase in AMI risk, and the cumulative time spent unemployed was also significantly associated with increased risk for AMI.⁵⁰ The risks remained significant after adjustment for socioeconomic, behavioral, and clinical variables.⁵⁰ The mechanisms underlying poorer outcomes with unemployment are not clear. In this study, time of entry into the workforce, which could suggest educational differences or health difficulties, was not significantly associated with outcomes.⁵⁰ A possible explanation is that of cumulative stress, because worse outcomes were seen with additional episodes of job loss and increased time spent unemployed. One limitation of the study was an older cohort (median age, 62 years) that may not be representative of the overall workforce.⁵⁰

ENVIRONMENTAL FACTORS

Neighborhood socioeconomic characteristics, in addition to individual SES, have been associated with CVD risk factors, adverse events, and mortality.^{15–17} A popu-

lation analysis of the Atherosclerosis Risk in Community study showed that living in disadvantaged areas is linked to higher incidence of CHD after controlling for individual income, education, and employment status.¹⁷ The findings were supported by a recent analysis of data from the Jackson Heart Study, which found an association between neighborhood disadvantage and cumulative biological risk, a score derived from biomarkers representing cardiovascular, metabolic, inflammatory, and neuroendocrine health.⁵¹ The environmental impact of health status and outcomes is driven by both physical and social attributes.^{16,52} Physical features of neighborhoods include the presence of sidewalks or recreational spaces, access to transportation, and availability and cost of healthy foods, whereas social attributes include safety, deprivation, social support, and lack of community cohesion.¹⁶ Socioeconomic differences in neighborhood characteristics can impact the availability of resources and influence promotion or maintenance of a healthy lifestyle. Favorable neighborhoods are associated with reduced cardiovascular risk factors, because long-term exposure to environments with greater access to physical activity and healthy food was associated with a lower incidence of diabetes mellitus and a lower prevalence of overweight or obesity.^{53,54} Other social characteristics, including neighborhood crime, also contribute to cardiovascular risk. In multiethnic populations, high individual- and neighborhood-level safety was associated with decreasing body mass index over time.⁵⁵ In contrast, poorly rated physical environments based on walking environment, availability of healthy food, safety, aesthetics, and social coherence were associated with elevated depressive symptoms and greater increase in waist circumference in individuals living in those neighborhoods.⁵⁶

Poor dietary options and the increased cost of healthy food may also contribute to the increased CVD risk of disadvantaged neighborhoods. Low-income areas have fewer food outlets and supermarkets, which results in limited access to fresh fruits and vegetables.^{57,58} The cost of healthy food is also higher in poor areas with limited access, in part because of the lower prevalence of supermarkets that typically offer lower prices and a variety of brand options.^{59,60} Low-income areas lacking access to healthy, affordable foods are known as food deserts, which are associated with diet-related chronic diseases and obesity.⁶¹ There are also racial disparities in food availability, because neighborhoods with a high prevalence of blacks tend to have more fast-food restaurants,⁵⁹ fewer supermarkets,⁶⁰ and fewer healthy options.⁶²

Neighborhood characteristics can change with time and provide additional insight into the relationship between environmental factors and cardiovascular health. Recent results from a longitudinal study with

a 12-year follow-up period showed that increasing the density of neighborhood healthy food resources was associated with lower coronary artery calcium over time. However, changes in other neighborhood characteristics, including walking and social environment, were not associated with changes in coronary artery calcium.⁶³ Further longitudinal studies have demonstrated that neighborhoods with favorable physical activity increased individual activities and improved cardiovascular risk factors.^{64,65} However, previous studies also showed that individuals often move into neighborhoods with similar SES across their life course, which impairs attempts to establish a causal relationship between environment and CVD.^{66,67} Unfortunately, a recent update by the US Department of Agriculture showed relatively small changes in low-income and low-access neighborhoods between 2010 and 2015 data.⁶⁸ Disadvantaged communities must be a priority to effectively reduce CVD disparities in individuals with low SES.

PSYCHOSOCIAL FACTORS AND CVD OUTCOMES

Psychosocial factors, including stress and depression, are strongly associated with adverse CVD outcomes, and growing evidence suggests they may disparately affect individuals of low SES.^{69,70} In analysis of the REGARDS study (Reasons for Geographic and Racial Differences in Stroke), a prospective observational cohort in the United States, individuals earning <\$35 000 annually who reported both stress and depressive symptoms had a 48% higher risk of developing CVD and a 33% increased risk of all-cause mortality after adjustment for socioeconomic, clinical, and behavioral factors.⁶⁹ The association was not present in subjects earning ≥\$35 000 per year.⁶⁹ Likewise, a study of psychological distress (depression, anxiety, and stress) demonstrated that the risk of CHD mortality was increased in those with low (hazard ratio, 1.31; $P=0.001$), but not high, SES ($P=0.107$).⁷⁰ The disparity in risk may relate to inadequate social or material resources to cope with stressful events and higher rates of adverse health behaviors, including smoking and physical inactivity, that are associated with stress and depression.^{71,72} The contribution of poor health behaviors in psychosocial distress is supported by a study from Ye et al⁷³ that found the increased risk of myocardial infarction or death in depressed individuals was attenuated after adjustment for behavioral mechanisms, in particular, smoking and physical inactivity. In addition to identification and treatment of the components of psychosocial stress, these findings suggest a role for aggressive targeting of smoking cessation and physical activity in these individuals.

SES AND CVD OUTCOMES BASED ON SEX

Women are overrepresented among those living in poverty; thus, they are disproportionately affected by the disparities in the distribution of wealth, income, and access to resources that ultimately can affect overall health and quality of life. In the WISE study (Women's Ischemic Syndrome Evaluation), women with ischemic heart disease in the lowest SES (household income <\$20 000/y) were more likely to be uninsured or have public insurance, yet had much higher drug costs and higher 5-year rehospitalization rates in comparison with higher-income women.⁷⁴ Income had a far greater impact than any other SES measure, including race, education, marital status, or employment status.⁷⁴ In a universal healthcare system (Southern Alberta, Canada), neighborhood SES was associated with the use of cardiac catheterization and 30-day mortality after acute coronary syndrome in women, but this same association was not seen in men.⁷⁵

INTERVENTIONS TO IMPROVE HEALTH BEHAVIORS AND RISK FACTORS

Programs aimed at improving health in the socially disadvantaged must first focus on aggressively targeting traditional risk factors that have strong associations with low SES.⁷ Behavioral counseling to reduce CVD risk factors including cholesterol levels, blood pressure, and diabetes mellitus incidence has been proven effective in the general population and is recommended by the US Preventive Services Task Force.^{76,77} Counseling of the family members of a patient with CVD, who are also at higher risk of vascular events, may improve dietary intake and physical activity, but further research into the applicability of these results to individuals with low SES is required. Studies have shown moderate success with smoking cessation programs aimed at patients of low SES.⁷⁸ Brown et al⁷⁸ used an Internet-based smoking cessation program to evaluate the success rate between socioeconomic groups and demonstrated a significant improvement in the low-SES cohort that was not realized in the high-SES population.

However, other studies have suggested more limited efficacy in individuals with low SES.^{79,80} Siren et al⁷⁹ found that behavioral counseling improved smoking cessation only in a high-education cohort and had no effect in a low-education group. In a study evaluating lifestyle interventions in low-income American Indians and Native Alaskans, the low-income groups had less improvement in body mass index, unhealthy food consumption, and physical activity in comparison with the high-income group.⁸⁰ The study did find an indirect relationship between the number of trained staff at each

location and the success of lifestyle modification.⁸⁰ Health counseling has been shown to improve diet in individuals with low SES, and interventions focused on the lack of convenient and affordable access to the components of a healthy diet may also be necessary to improve dietary habits in low-SES areas.⁷⁹

Regular physical activity is associated with a decreased risk of CVD and its comorbidities.^{81,82} Likewise, physical inactivity is an independent risk factor for poor CVD outcomes.⁸³ However, despite the known impact of physical activity and inactivity on CVD modulation, fewer than half of US adults met the recommended levels of physical activity in 2011.⁸⁴ The burden of inadequate leisure-time physical activity is especially pronounced in individuals of low SES, which may be because of increased occupational responsibilities or reduced access to safe facilities for exercise.^{85,86} Programs aimed at increasing physical activity have demonstrated promising results in individuals with low SES.^{79,87} The Walk Your Heart to Health program successfully improved physical activity in a low- to middle-income group consisting primarily of ethnic minorities in Detroit, Michigan.⁸⁷ The intervention group met 3 times weekly at community sites and walked for an increasing amount of time (45–90 minutes). Physical activity of the group was increased at 8 weeks, and improvements were noted in the cohort's systolic blood pressure, fasting blood glucose, total cholesterol, waist circumference, and body mass index at 8 weeks that were maintained at 32 weeks.⁸⁷ Physical activity appears to be an important target in low-SES communities because of its widespread benefits and inherently inexpensive nature; however, physical insecurity (violence, crime) or lack of infrastructure (sidewalks, bicycle paths) in the neighborhood social environment may severely impact an individual's ability to create a sustainable physical activity regimen.⁸⁸ Thus, policy intervention to address such factors in low-SES populations must be addressed in conjunction with efforts to change individual behavior.⁸⁹

A substantial barrier to the primary prevention of CVD, especially in low-SES groups, is a lack of access to healthcare providers. Given the increasing demands on physicians, strategies have been developed to better appropriate available resources. One effective community-based strategy using this paradigm is task shifting, an idea conceived by the World Health Organization as early as 1980 and defined as redistributing primary care responsibilities from physician to nonphysician providers.⁹⁰ Although a dearth of evidence exists for the use of task shifting in HIC, the trial by Adeyemo et al⁹¹ in Nigeria, a LMIC, demonstrated a reduction in systolic blood pressure from a preintervention mean of 168/92 to a goal of <140/90 in 66.7% of patients by using nurse-driven task shifting in comparison with 65.4% of patients receiving usual medical care. Task shifting can

be further extended to include health workers without formal healthcare training, known as community health workers. Community health workers play a valuable role in cost-effective screening for CVD in low-resource countries and communities.^{92,93} Given the constraints of modern medical practice, task shifting may have utility for CVD screening and risk factor improvement in areas with reduced resources.

Numerous studies have evaluated the role of community interventions on major adverse cardiovascular events and risk factors. Mass media negatively portraying tobacco reduces smoking prevalence by ≈2%.⁹⁴ Tobacco cessation rates can be further impacted by increasing prices of tobacco products and state programs to provide nicotine replacement therapy.⁹⁴ Task shifting the role of smoking cessation management to a community pharmacist increased the quit rate from 2.7% to 14.3% in study participants.⁹⁴ Similar effects are seen with mass media advocacy for salt restriction and dietary modification.⁹⁴ Subsidization of healthy foods is another community-level intervention that has demonstrated benefit in low-income individuals. Reducing the cost of healthy foods has been shown to improve diets and reduce the barriers to a healthy diet in food deserts.⁹⁵ Similarly, taxation of unhealthy foods decreases their consumption.⁹⁵ A combined strategy can be used to improve diets in at-risk communities. Environmental engineering, including construction of new supermarkets to increase access to healthy foods, still lacks adequate data to support its use.⁹⁵

Heart disease exhibits regional clustering in the United States (Figure 2, map created using the Interactive Atlas of Heart Disease and Stroke).⁹⁶ Geomapping, the identification of geographic hot spots of individuals at high risk for CVD, may be used in the future to target communities that would benefit from aggressive community-based interventions.⁹⁷ Geomapping has demonstrated promising early results in Sweden for identification of at-risk populations for diabetes mellitus and could be adapted to provide increased screening and treatment services for low-SES communities with a high prevalence of CVD risk.⁹⁸

Strategies for identifying and improving health disparities in the United States were addressed in a Think Tank meeting organized by The National Heart, Lung, and Blood Institute.⁹⁹ The panel recommendations focus on the conversion of evidence to implementation through an increased breadth of research themes and transdisciplinary training programs to expand research capacities.⁹⁹ The panel further emphasized the role of platforms to optimize research and the development of collaborator and stakeholder networks to create benchmark studies.⁹⁹ See Figure 3 for a summary of potential interventions in individuals with low SES.

Targeting lower socioeconomic populations certainly must involve multilevel behavioral interventions at the




Figure 2. Cardiovascular disease death rates in individuals age ≥ 35 years by county between 2013 and 2015.

Rates are spatially smoothed to enhance the stability of rates in counties with small populations. Data sources are the National Vital Statistics System and National Center for Health Statistics. This map was created using the Interactive Atlas of Heart Disease and Stroke, a website developed by the Centers for Disease Control and Prevention, Division for Heart Disease and Stroke Prevention.⁹⁶

individual and community levels that incorporate a comprehensive understanding of social determinants of health and a concerted effort to engage the community.⁸⁹ Specific efforts to reform the socioecological environment for primordial and primary prevention, as outlined by the “American Heart Association Guide for Improving Cardiovascular Health at the Community Level,” include an emphasis to direct resources to support legislation encouraging smoking cessation, promote sodium restriction in processed foods, and sup-

port modifications in the environment to encourage physical activity.⁸⁹ Through the use of existing cohesive networks such as healthcare facilities, schools, religious organizations, worksites, social networks, media networks, and virtual communities, physicians, policy makers, and activists can develop a more effective platform to engage and intervene in communities at a grassroots level.⁸⁹ Unfortunately, data addressing improvement in CVD in lower-SES populations through targeted intervention are limited, and further studies should be




Figure 3. Association between low socioeconomic status, cardiovascular risk factors, cardiovascular disease, and interventions. CVD indicates cardiovascular disease; and SES, socioeconomic status.

performed to investigate the potential impact of such interventions.⁸⁸ In addition, although community interventions play an essential role in changing individual behavior, macrolevel factors may significantly impede the capacity to change individual behavior and should also be addressed.

FUTURE TRENDS AND DIRECTIONS

A study in England determined that CHD mortality was decreasing in individuals of all SES, but the rate of decline was steepest in the most affluent group in comparison with those with lower SES.¹⁰⁰ Lotufo et al¹⁰¹ noted similar findings in a comparison of high- and low-income groups in Brazil, a high-middle income country. Several countries have also seen flattening of the reductions in CHD mortality in younger individuals of low SES.^{102,103} In the United States and Scotland, the trend may be linked to increasing inequalities in smoking in low socioeconomic groups.^{102,103} The United States has also experienced an increase in diabetes mellitus among the socially deprived.^{102,104} These trends are concerning and suggest that, although overall cardiovascular care is improving, the advances are preferentially helping the socially advantaged and widening the gap of health inequality.

Properly risk-stratifying patients will be a critical aspect of identifying patients with low SES with an increased risk of CVD that would not be conveyed by traditional risk factors alone. The ASSIGN score (Assessing Cardiovascular Risk Using Scottish Intercollegiate Guidelines Network Guidelines) and QRISK algorithm in the United Kingdom are alternative risk stratification tools that integrate postal code income with traditional CVD risk factors.^{105,106} The ASSIGN score provided a statistically significant, but marginal, improvement in risk prediction in comparison with the Framingham Risk Score (FRS).¹⁰⁵ The QRISK algorithm also improved prediction of CVD events in individuals whose risk was underpredicted by the FRS.¹⁰⁶ Although QRISK and ASSIGN are meant to serve as alternative approaches to the FRS, Franks et al¹⁰⁷ incorporated SES markers directly into the existing FRS framework. The integration of educational attainment and income as additional markers to predict risk into the FRS removed the SES bias seen in individuals with low SES.¹⁰⁷ Accurate prediction is important to identify appropriate candidates for aggressive primary prevention, which includes statin therapy in individuals at borderline risk for CVD or additional resources dedicated to counseling in patients with multiple markers for low SES. Models incorporating SES for risk prediction are challenging because of the regional and cultural differences in SES markers. It remains to be seen if a single risk stratification system will be adequate or if regional variants will be required, particularly

with the use of the American College of Cardiology and American Heart Association atherosclerotic CVD risk assessment tool in the US population.

Several organizations have created initiatives to reduce health disparities in the United States and worldwide. In 2011, the World Health Organization set a goal to reduce premature deaths from noncommunicable diseases, including CVD, by 25% by 2025 (25by25).¹⁰⁸ The World Health Organization action plan targets modifiable CVD risk factors: tobacco use, alcohol intake, sodium intake, physical inactivity, diabetes mellitus/obesity, and hypertension.¹⁰⁹ The World Heart Federation expanded that goal to include a 25% reduction in CVD mortality by the year 2025 through 9 strategies, including reduction in tobacco use, detection and treatment of hypertension, improved access to proven CVD medications, implementation of community-based interventions, creation of large studies to characterize lifestyle habits of regional populations, and the development of partnerships between high- and low-resource countries to facilitate the transfer of knowledge and funding.^{110,111}

Reducing disparities in health will require a multilevel, collaborative approach. The overarching principles include a focus on identifying individuals and communities at greatest risk and putting more resources toward these groups, improving access to quality health care, increasing cultural competence, and revamping medical education. Government (federal and local), businesses/employers, healthcare systems, schools, community organizations, and individuals/families all have an integral role in elimination of health disparities (Figure 4). Furthermore, because the foundation of SES and health in adulthood are influenced by conditions in childhood, targeted preschool and early childhood interventions have important implications for reducing disparities.¹¹²




Figure 4. Interventions at the community, government, healthcare systems, and individual levels should be targeted equally in efforts to reduce disparities in health.

The focus of future studies should be on the implementation of research evaluating strategies to reduce health disparities. These studies should be designed to identify effective policy changes and program interventions to reduce disparities. Furthermore, the costs and benefits of individual- and community-level interventions to identify the most cost-effective strategies and interventions should be studied.¹¹³

CONCLUSIONS

SES has a measurable and significant impact on cardiovascular health. Individuals of low SES carry a substantial burden of CVD and are more likely to experience increased event rates and poorer outcomes. Current models do not adequately account for the risk conveyed by low SES. We now have compelling evidence showing that the independent association between SES and mortality is comparable in strength and consistency to that of the traditional major risk factors.³ As Tobias concluded, the strength of that evidence “is now impossible to ignore.”¹¹⁴ The time has come for increased focus on effective and sustainable interventions informed by clinical and population science insights from SES research. In addition, further research is required to better understand the underlying mechanisms of CVD risk that disproportionately affect individuals of low SES. Once the causes of the discrepancies in health equity are better understood, targeted interventions can be pursued to better address disparities in populations at risk.

ARTICLE INFORMATION

Correspondence

Laurence S. Sperling, MD, Division of Cardiology, Department of Medicine, Emory University, 1365 Clifton Rd NE, Building A, Suite 2200, Atlanta, GA 30322. E-mail lsperli@emory.edu

Affiliations

Department of Medicine (W.M.S., J.C.L., T.V.), Emory Clinical Cardiovascular Research Institute (H.M.K., J.S., P.S., A.A.Q., L.S.S.), Emory University School of Medicine, Atlanta, GA. Morehouse School of Medicine, Atlanta, GA (H.A.T.). University of Arizona-Phoenix College of Medicine (M.G.). Cedars-Sinai Heart Institute, Cedars-Sinai Medical Center, Los Angeles, CA (J.G.H.). Hofstra Northwell School of Medicine, Hempstead, NY (J.H.M.). Tulane University School of Medicine, New Orleans, LA (K.C.F.). Center for Translation Research and Implementation Science, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD (G.A.M.).

Sources of Funding

Dr Sandesara is supported by the Abraham J. & Phyllis Katz Foundation (Atlanta, GA). Dr Quyyumi is supported by the National Institutes of Health (1R61HL138657-01, 5P01HL101398-02, 1P20HL113451-01, 1P30DK111024-01, 1RF1AG051633-01, R01 NS064162-01, R01 HL89650-01, HL095479-01, 1U10HL110302-01, 1DP3DK094346-01, and 2P01HL086773-06A1) and the American Heart Association (AHA 0000031288).

Disclosures

None.

REFERENCES

- Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, de Ferranti SD, Floyd J, Fornage M, Gillespie C, Isasi CR, Jiménez MC, Jordan LC, Judd SE, Lackland D, Lichtman JH, Lisabeth L, Liu S, Longenecker CT, Mackey RH, Matsushita K, Mozaffarian D, Mussolino ME, Nasir K, Neumar RW, Palaniappan L, Pandey DK, Thiagarajan RR, Reeves MJ, Ritchey M, Rodriguez CJ, Roth GA, Rosamond WD, Sasson C, Towfighi A, Tsao CW, Turner MB, Virani SS, Voeks JH, Willey JZ, Wilkins JT, Wu JH, Alger HM, Wong SS, Muntner P; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2017 Update: a report from the American Heart Association. *Circulation*. 2017;135:e146–e603. doi: 10.1161/CIR.0000000000000485.
- Franks P, Winters PC, Tancredi DJ, Fiscella KA. Do changes in traditional coronary heart disease risk factors over time explain the association between socio-economic status and coronary heart disease? *BMC Cardiovasc Disord*. 2011;11:28. doi: 10.1186/1471-2261-11-28.
- Stringhini S, Carmeli C, Jokela M, Avendaño M, Muennig P, Guida F, Ricceri F, d'Errico A, Barros H, Bochud M, Chadeau-Hyam M, Clavel-Chapelon F, Costa G, Delpierre C, Fraga S, Goldberg M, Giles GG, Krogh V, Kelly-Irving M, Layte R, Lasserre AM, Marmot MG, Preisig M, Shipley MJ, Vollenweider P, Zins M, Kawachi I, Steptoe A, Mackenbach JP, Vineis P, Kivimäki M; LIFEPAth consortium. Socioeconomic status and the 25 × 25 risk factors as determinants of premature mortality: a multicohort study and meta-analysis of 1.7 million men and women. *Lancet*. 2017;389:1229–1237. doi: 10.1016/S0140-6736(16)32380-7.
- Lynch JW, Kaplan GA, Cohen RD, Tuomilehto J, Salonen JT. Do cardiovascular risk factors explain the relation between socioeconomic status, risk of all-cause mortality, cardiovascular mortality, and acute myocardial infarction? *Am J Epidemiol*. 1996;144:934–942.
- Winkleby MA, Jatulis DE, Frank E, Fortmann SP. Socioeconomic status and health: how education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Public Health*. 1992;82:816–820.
- Adler NE, Glymour MM, Fielding J. Addressing social determinants of health and health inequalities. *JAMA*. 2016;316:1641–1642. doi: 10.1001/jama.2016.14058.
- Davis AM, Vinci LM, Okwuosa TM, Chase AR, Huang ES. Cardiovascular health disparities: a systematic review of health care interventions. *Med Care Res Rev*. 2007;64(5 suppl):295–300S. doi: 10.1177/1077558707305416.
- Mackenbach JP, Cavelaars AE, Kunst AE, Groenou F. Socioeconomic inequalities in cardiovascular disease mortality: an international study. *Eur Heart J*. 2000;21:1141–1151. doi: 10.1053/euhj.1999.1990.
- Gebreab SY, Diez Roux AV, Brenner AB, Hickson DA, Sims M, Subramanyam M, Griswold ME, Wyatt SB, James SA. The impact of life-course socioeconomic position on cardiovascular disease events in African Americans: the Jackson Heart Study. *J Am Heart Assoc*. 2015;4:e001553. doi: 10.1161/JAHA.114.001553.
- Janković J, Erić M, Stojisavljević D, Marinković J, Janković S. Socio-economic differences in cardiovascular health: findings from a cross-sectional study in a middle-income country. *PLoS One*. 2015;10:e0141731. doi: 10.1371/journal.pone.0141731.
- Mosquera PA, San Sebastian M, Waenerlund AK, Ivarsson A, Weinehall L, Gustafsson PE. Income-related inequalities in cardiovascular disease from mid-life to old age in a Northern Swedish cohort: a decomposition analysis. *Soc Sci Med*. 2016;149:135–144. doi: 10.1016/j.socscimed.2015.12.017.
- Kucharska-Newton AM, Harald K, Rosamond WD, Rose KM, Rea TD, Salomaa V. Socioeconomic indicators and the risk of acute coronary heart disease events: comparison of population-based data from the United States and Finland. *Ann Epidemiol*. 2011;21:572–579. doi: 10.1016/j.annepidem.2011.04.006.
- Liu K, Cedres LB, Stamler J, Dyer A, Stamler R, Nanas S, Berkson DM, Paul O, Lepper M, Lindberg HA, Marquardt J, Stevens E, Schoenberger JA, Shekelle RB, Collette P, Shekelle S, Garside D. Relationship of education to major risk factors and death from coronary heart disease, cardiovascular diseases and all causes, findings of three Chicago epidemiologic studies. *Circulation*. 1982;66:1308–1314.
- Meneton P, Kesse-Guyot E, Méjean C, Fezeu L, Galan P, Hercberg S, Ménéard J. Unemployment is associated with high cardiovascular event rate and increased all-cause mortality in middle-aged socially privileged individuals. *Int Arch Occup Environ Health*. 2015;88:707–716. doi: 10.1007/s00420-014-0997-7.
- Clark CR, Ommerborn MJ, Hickson DA, Grooms KN, Sims M, Taylor HA, Albert MA. Neighborhood disadvantage, neighborhood safety and cardiometabolic risk factors in African Americans: biosocial associations

- in the Jackson Heart study. *PLoS One*. 2013;8:e63254. doi: 10.1371/journal.pone.0063254.
16. Diez Roux AV. Residential environments and cardiovascular risk. *J Urban Health*. 2003;80:569–589. doi: 10.1093/jurban/jtg065.
 17. Diez Roux AV, Merkin SS, Arnett D, Chambless L, Massing M, Nieto FJ, Sorlie P, Szklo M, Tyroler HA, Watson RL. Neighborhood of residence and incidence of coronary heart disease. *N Engl J Med*. 2001;345:99–106. doi: 10.1056/NEJM200107123450205.
 18. Lewis MW, Khodneva Y, Redmond N, Durant RW, Judd SE, Wilkinson LL, Howard VJ, Safford MM. The impact of the combination of income and education on the incidence of coronary heart disease in the prospective Reasons for Geographic and Racial Differences in Stroke (REGARDS) cohort study. *BMC Public Health*. 2015;15:1312. doi: 10.1186/s12889-015-2630-4.
 19. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, Ezzati M, Shibuya K, Salomon JA, Abdalla S, Aboyans V, Abraham J, Ackerman I, Aggarwal R, Ahn SY, Ali MK, Alvarado M, Anderson HR, Anderson LM, Andrews KG, Atkinson C, Baddour LM, Bahalim AN, Barker-Collo S, Barrero LH, Bartels DH, Basáñez MG, Baxter A, Bell ML, Benjamin EJ, Bennett D, Bernabé E, Bhalla K, Bhandari B, Bikbov B, Bin Abdulhak A, Birbeck G, Black JA, Blencowe H, Blore JD, Blyth F, Bolliger I, Bonaventure A, Boufous S, Bourne R, Boussinesq M, Braithwaite T, Brayne C, Bridgett L, Brooker S, Brooks P, Brugha TS, Bryan-Hancock C, Bucello C, Buchbinder R, Buckle G, Budke CM, Burch M, Burney P, Burstein R, Calabria B, Campbell B, Canter CE, Carabin H, Carapetis J, Carmona L, Cella C, Charlson F, Chen H, Cheng AT, Chou D, Chugh SS, Coffeng LE, Colan SD, Colquhoun S, Colson KE, Condon J, Connor MD, Cooper LT, Corriere M, Cortinovis M, de Vaccaro KC, Couser W, Cowie BC, Criqui MH, Cross M, Dabhadkar KC, Dahiya M, Dahodwala N, Damsere-Derry J, Danaei G, Davis A, De Leo D, Degenhardt L, Dellavalle R, Delossantos A, Denenberg J, Derrett S, Des Jarlais DC, Dharmaratne SD, Dherani M, Diaz-Torne C, Dolk H, Dorsey ER, Driscoll T, Duber H, Ebel B, Edmond K, Elbaz A, Ali SE, Erskine H, Erwin PJ, Espindola P, Ewoigbokhan SE, Farzadfar F, Feigin V, Felson DT, Ferrari A, Ferri CP, Fèvre EM, Finucane MM, Flaxman S, Flood L, Foreman K, Forouzanfar MH, Fowkes FG, Fransen M, Freeman MK, Gabbe BJ, Gabriel SE, Gakidou E, Ganatra HA, Garcia B, Gaspari F, Gillum RF, Gmel G, Gonzalez-Medina D, Gosselin R, Grainger R, Grant B, Groeger J, Guillemin F, Gunnell D, Gupta R, Haagsma J, Hagan H, Halasa YA, Hall W, Haring D, Haro JM, Harrison JE, Havmoeller R, Hay RJ, Higashi H, Hill C, Hoen B, Hoffman H, Hotez PJ, Hoy D, Huang JJ, Ibeanusi SE, Jacobsen KH, James SL, Jarvis D, Jasrasaria R, Jayaraman S, Johns N, Jonas JB, Karthikeyan G, Kassebaum N, Kawakami N, Keren A, Khoo JP, King CH, Knowlton LM, Kobusingye O, Koranteng A, Krishnamurthi R, Laden F, Laloo R, Laslett LL, Lathlean T, Leasher JL, Lee YY, Leigh J, Levinson D, Lim SS, Limb E, Lin JK, Lipnick M, Lipshultz SE, Liu W, Loane M, Ohno SL, Lyons R, Mabwajano J, MacIntyre MF, Malekzadeh R, Mallinger L, Manivannan S, Marcenies W, March L, Margolis DJ, Marks GB, Marks R, Matsumori A, Matzopoulos R, Mayosi BM, McAnulty JH, McDermott MM, McGill N, McGrath J, Medina-Mora ME, Meltzer M, Mensah GA, Merriman TR, Meyer AC, Miglioli V, Miller M, Miller TR, Mitchell PB, Mock C, Mocumbi AO, Moffitt TE, Mokdad AA, Monasta L, Montico M, Moradi-Lakeh M, Moran A, Morawska L, Mori R, Murdoch ME, Mwaniki MK, Naidoo K, Nair MN, Naldi L, Narayan KM, Nelson PK, Nelson RG, Nevitt MC, Newton CR, Nolte S, Norman P, Norman R, O'Donnell M, O'Hanlon S, Olives C, Omer SB, Ortblad K, Osborne R, Ozgediz D, Page A, Pahari B, Pandian JD, Rivero AP, Patten SB, Pearce N, Padilla RP, Perez-Ruiz F, Perico N, Pesudovs K, Phillips D, Phillips MR, Pierce K, Pion S, Polanczyk GV, Polinder S, Pope CA 3rd, Popova S, Porrini E, Pourmalek F, Prince M, Pullan RL, Ramaiah KD, Ranganathan D, Razavi H, Regan M, Rehm JT, Rein DB, Remuzzi G, Richardson K, Rivara FP, Roberts T, Robinson C, De León FR, Ronfani L, Room R, Rosenfeld LC, Rushton L, Sacco RL, Saha S, Sampson U, Sanchez-Riera L, Sanman E, Schwebel DC, Scott JG, Segui-Gomez M, Shahraz S, Shepard DS, Shin H, Shivakoti R, Singh D, Singh GM, Singh JA, Singleton J, Sleet DA, Sliwa K, Smith E, Smith JL, Stapelberg NJ, Steer A, Steiner T, Stolk WA, Stovner LJ, Sudfeld C, Syed S, Tamburlini G, Tavakkoli M, Taylor HR, Taylor JA, Taylor WJ, Thomas B, Thomson WM, Thurston GD, Tleyeh IM, Tonelli M, Towbin JA, Truelsén T, Tsilimbaris MK, Ubeda C, Undurraga EA, van der Werf MJ, van Os J, Vavilala MS, Venketasubramanian N, Wang M, Wang W, Watt K, Weatherall DJ, Weinstock MA, Weintraub R, Weisskopf MG, Weissman MM, White RA, Whiteford H, Wiebe N, Wiersma ST, Wilkinson JD, Williams HC, Williams SR, Witt E, Wolfe F, Woolf AD, Wulf S, Yeh PH, Zaidi AK, Zheng ZJ, Zonies D, Lopez AD, AlMazroa MA, Memish ZA. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2197–2223. doi: 10.1016/S0140-6736(12)61689-4.
 20. Goyal A, Bhatt DL, Steg PG, Gersh BJ, Alberts MJ, Ohman EM, Corbaccán R, Eagle KA, Gaxiola E, Gao R, Goto S, D'Agostino RB, Califf RM, Smith SC Jr, Wilson PW; Reduction of Atherothrombosis for Continued Health (REACH) Registry Investigators. Attained educational level and incident atherothrombotic events in low- and middle-income compared with high-income countries. *Circulation*. 2010;122:1167–1175. doi: 10.1161/CIRCULATIONAHA.109.919274.
 21. Dinsa GD, Goryakin Y, Fumagalli E, Suhrcke M. Obesity and socioeconomic status in developing countries: a systematic review. *Obes Rev*. 2012;13:1067–1079. doi: 10.1111/j.1467-789X.2012.01017.x.
 22. Gerber Y, Weston SA, Killian JM, Therneau TM, Jacobsen SJ, Roger VL. Neighborhood income and individual education: effect on survival after myocardial infarction. *Mayo Clin Proc*. 2008;83:663–669. doi: 10.4065/83.6.663.
 23. Stirbu I, Looman C, Nijhof GJ, Reulings PG, Mackenbach JP. Income inequalities in case death of ischaemic heart disease in the Netherlands: a national record-linked study. *J Epidemiol Community Health*. 2012;66:1159–1166. doi: 10.1136/jech-2011-200924.
 24. Yong CM, Abnoui F, Asch SM, Heidenreich PA. Socioeconomic inequalities in quality of care and outcomes among patients with acute coronary syndrome in the modern era of drug eluting stents. *J Am Heart Assoc*. 2014;3:e001029. doi: 10.1161/JAHA.14.001029.
 25. Wang JY, Wang CY, Juang SY, Huang KY, Chou P, Chen CW, Lee CC. Low socioeconomic status increases short-term mortality of acute myocardial infarction despite universal health coverage. *Int J Cardiol*. 2014;172:82–87. doi: 10.1016/j.ijcard.2013.12.082.
 26. Dunlay SM, Pack QR, Thomas RJ, Killian JM, Roger VL. Participation in cardiac rehabilitation, readmissions, and death after acute myocardial infarction. *Am J Med*. 2014;127:538–546. doi: 10.1016/j.amjmed.2014.02.008.
 27. Doll JA, Hellkamp A, Thomas L, Ho PM, Kontos MC, Whoolley MA, Boyden TF, Peterson ED, Wang TY. Effectiveness of cardiac rehabilitation among older patients after acute myocardial infarction. *Am Heart J*. 2015;170:855–864.
 28. Lemstra ME, Alsabbagh W, Rajakumar RJ, Rogers MR, Blackburn D. Neighborhood income and cardiac rehabilitation access as determinants of nonattendance and noncompletion. *Can J Cardiol*. 2013;29:1599–1603.
 29. Rasmussen JN, Gislason GH, Rasmussen S, Abildstrom SZ, Schramm TK, Køber L, Diderichsen F, Osler M, Torp-Pedersen C, Madsen M. Use of statins and beta-blockers after acute myocardial infarction according to income and education. *J Epidemiol Community Health*. 2007;61:1091–1097. doi: 10.1136/jech.2006.055525.
 30. Hanley GE, Morgan S, Reid RJ. Income-related inequity in initiation of evidence-based therapies among patients with acute myocardial infarction. *J Gen Intern Med*. 2011;26:1329–1335. doi: 10.1007/s11606-011-1799-1.
 31. Yu TH, Hou YC, Chung KP. Do low-income coronary artery bypass surgery patients have equal opportunity to access excellent quality of care and enjoy good outcome in Taiwan? *Int J Equity Health*. 2014;13:64. doi: 10.1186/s12939-014-0064-8.
 32. Parikh PB, Yang J, Leigh S, Dorjee K, Parikh R, Sakellarios N, Meng H, Brown DL. The impact of financial barriers on access to care, quality of care and vascular morbidity among patients with diabetes and coronary heart disease. *J Gen Intern Med*. 2014;29:76–81. doi: 10.1007/s11606-013-2635-6.
 33. Joynt KE, Orav EJ, Jha AK. Thirty-day readmission rates for Medicare beneficiaries by race and site of care. *JAMA*. 2011;305:675–681. doi: 10.1001/jama.2011.123.
 34. Gusmano MK, Weisz D, Rodwin VG, Lang J, Qian M, Bocquier A, Moysan V, Verger P. Disparities in access to health care in three French regions. *Health Policy*. 2014;114:31–40. doi: 10.1016/j.healthpol.2013.07.011.
 35. Woodward M, Peters SA, Batty GD, Ueshima H, Woo J, Giles GG, Barzi F, Ho SC, Huxley RR, Arima H, Fang X, Dobson A, Lam TH, Vathesatogkit P; Asia Pacific Cohort Studies Collaboration. Socioeconomic status in relation to cardiovascular disease and cause-specific mortality: a comparison of Asian and Australasian populations in a pooled analysis. *BMJ Open*. 2015;5:e006408. doi: 10.1136/bmjopen-2014-006408.
 36. Hu B, Li W, Wang X, Liu L, Teo K, Yusuf S; INTER-HEART Investigators. Marital status, education, and risk of acute myocardial infarction in Mainland China: the INTER-HEART study. *J Epidemiol*. 2012;22:123–129.
 37. Kelly MJ, Weitzen S. The association of lifetime education with the prevalence of myocardial infarction: an analysis of the 2006 Behavioral Risk Factor Surveillance System. *J Community Health*. 2010;35:76–80. doi: 10.1007/s10900-009-9189-x.

38. Rosengren A, Subramanian SV, Islam S, Chow CK, Avezum A, Kazmi K, Sliwa K, Zubaid M, Rangarajan S, Yusuf S; INTERHEART Investigators. Education and risk for acute myocardial infarction in 52 high, middle and low-income countries: INTERHEART case-control study. *Heart*. 2009;95:2014–2022. doi: 10.1136/hrt.2009.182436.
39. Mehta RH, O'Shea JC, Stebbins AL, Granger CB, Armstrong PW, White HD, Topol EJ, Califf RM, Ohman EM. Association of mortality with years of education in patients with ST-segment elevation myocardial infarction treated with fibrinolysis. *J Am Coll Cardiol*. 2011;57:138–146. doi: 10.1016/j.jacc.2010.09.021.
40. Gerber Y, Goldbourt U, Drory Y; Israel Study Group on First Acute Myocardial Infarction. Interaction between income and education in predicting long-term survival after acute myocardial infarction. *Eur J Cardiovasc Prev Rehabil*. 2008;15:526–532. doi: 10.1097/HJR.0b013e328304feac.
41. Kim JH, Jeong MH, Park IH, Choi JS, Rhee JA, Lee DH, Park SH, Kim IS, Jeong HC, Cho JY, Jang SY, Lee KH, Park KH, Sim DS, Kim KH, Hong YJ, Park HW, Kim JH, Ahn Y, Cho JG, Park JC. The association of socioeconomic status with three-year clinical outcomes in patients with acute myocardial infarction who underwent percutaneous coronary intervention. *J Korean Med Sci*. 2014;29:536–543. doi: 10.3346/jkms.2014.29.4.536.
42. Alter DA, Iron K, Austin PC, Naylor CD; SESAMI Study Group. Socioeconomic status, service patterns, and perceptions of care among survivors of acute myocardial infarction in Canada. *JAMA*. 2004;291:1100–1107. doi: 10.1001/jama.291.9.1100.
43. Kershaw KN, Droomers M, Robinson WR, Carnethon MR, Daviglus ML, Monique Verschuren WM. Quantifying the contributions of behavioral and biological risk factors to socioeconomic disparities in coronary heart disease incidence: the MORGEN study. *Eur J Epidemiol*. 2013;28:807–814. doi: 10.1007/s10654-013-9847-2.
44. Luepker RV, Rosamond WD, Murphy R, Sprafka JM, Folsom AR, McGovern PG, Blackburn H. Socioeconomic status and coronary heart disease risk factor trends. The Minnesota Heart Survey. *Circulation*. 1993;88(5 pt 1):2172–2179.
45. Cajita MI, Cajita TR, Han HR. Health literacy and heart failure: a systematic review. *J Cardiovasc Nurs*. 2016;31:121–130. doi: 10.1097/JCN.0000000000000229.
46. Zhang NJ, Terry A, McHorney CA. Impact of health literacy on medication adherence: a systematic review and meta-analysis. *Ann Pharmacother*. 2014;48:741–751. doi: 10.1177/1060028014526562.
47. Bostock S, Steptoe A. Association between low functional health literacy and mortality in older adults: longitudinal cohort study. *BMJ*. 2012;344:e1602.
48. Loucks EB, Gilman SE, Howe CJ, Kawachi I, Kubzansky LD, Rudd RE, Martin LT, Nandi A, Wilhelm A, Buka SL. Education and coronary heart disease risk: potential mechanisms such as literacy, perceived constraints, and depressive symptoms. *Health Educ Behav*. 2015;42:370–379. doi: 10.1177/1090198114560020.
49. Méjean C, Droomers M, van der Schouw YT, Sluijs I, Czernichow S, Grobbee DE, Bueno-de-Mesquita HB, Beulens JW. The contribution of diet and lifestyle to socioeconomic inequalities in cardiovascular morbidity and mortality. *Int J Cardiol*. 2013;168:5190–5195. doi: 10.1016/j.ijcard.2013.07.188.
50. Dupre ME, George LK, Liu G, Peterson ED. The cumulative effect of unemployment on risks for acute myocardial infarction. *Arch Intern Med*. 2012;172:1731–1737.
51. Barber S, Hickson DA, Kawachi I, Subramanian SV, Earls F. Neighborhood disadvantage and cumulative biological risk among a socioeconomically diverse sample of African American adults: an examination in the Jackson Heart Study. *J Racial Ethn Health Disparities*. 2016;3:444–456. doi: 10.1007/s40615-015-0157-0.
52. Diez-Roux AV, Nieto FJ, Muntaner C, Tyroler HA, Comstock GW, Shahar E, Cooper LS, Watson RL, Szklo M. Neighborhood environments and coronary heart disease: a multilevel analysis. *Am J Epidemiol*. 1997;146:48–63.
53. Christine PJ, Auchincloss AH, Bertoni AG, Carnethon MR, Sánchez BN, Moore K, Adar SD, Horwich TB, Watson KE, Diez Roux AV. Longitudinal associations between neighborhood physical and social environments and incident type 2 diabetes mellitus: the Multi-Ethnic Study of Atherosclerosis (MESA). *JAMA Intern Med*. 2015;175:1311–1320. doi: 10.1001/jamainternmed.2015.2691.
54. Creatore MI, Glazier RH, Moineddin R, Fazli GS, Johns A, Gozdyra P, Matheison FI, Kaufman-Shriqui V, Rosella LC, Manuel DG, Booth GL. Association of neighborhood walkability with change in overweight, obesity, and diabetes. *JAMA*. 2016;315:2211–2220. doi: 10.1001/jama.2016.5898.
55. Powell-Wiley TM, Moore K, Allen N, Block R, Evenson KR, Mujahid M, Diez Roux AV. Associations of neighborhood crime and safety and with changes in body mass index and waist circumference: the Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol*. 2017;186:280–288. doi: 10.1093/aje/kwx082.
56. Remigio-Baker RA, Diez Roux AV, Szklo M, Crum RM, Leoutsakos JM, Franco M, Schreiner PJ, Carnethon MR, Nettleton JA, Mujahid MS, Michos ED, Gary-Webb TL, Golden SH. Physical environment may modify the association between depressive symptoms and change in waist circumference: the Multi-Ethnic Study of Atherosclerosis. *Psychosomatics*. 2014;55:144–154. doi: 10.1016/j.psych.2013.10.008.
57. Alwitt LF, Donley TD. Retail stores in poor urban neighborhoods. *J Consum Aff*. 1997;139–164.
58. Giang T, Karpyn A, Laurison HB, Hillier A, Perry RD. Closing the grocery gap in underserved communities: the creation of the Pennsylvania Fresh Food Financing Initiative. *J Public Health Manag Pract*. 2008;14:272–279. doi: 10.1097/01.PHH.0000316486.57512.bf.
59. Block JP, Scribner RA, DeSalvo KB. Fast food, race/ethnicity, and income: a geographic analysis. *Am J Prev Med*. 2004;27:211–217. doi: 10.1016/j.amepre.2004.06.007.
60. Block D, Kouba J. A comparison of the availability and affordability of a market basket in two communities in the Chicago area. *Public Health Nutr*. 2006;9:837–845.
61. Dubowitz T, Zenk SN, Ghosh-Dastidar B, Cohen DA, Beckman R, Hunter G, Steiner ED, Collins RL. Healthy food access for urban food desert residents: examination of the food environment, food purchasing practices, diet and BMI. *Public Health Nutr*. 2015;18:2220–2230.
62. Lewis LB, Sloane DC, Nascimento LM, Diamant AL, Guinyard JJ, Yancey AK, Flynn G; REACH Coalition of the African Americans Building a Legacy of Health Project. African Americans' access to healthy food options in South Los Angeles restaurants. *Am J Public Health*. 2005;95:668–673. doi: 10.2105/AJPH.2004.050260.
63. Wing JJ, August E, Adar SD, Dannenberg AL, Hajat A, Sánchez BN, Stein JH, Tattersall MC, Diez Roux AV. Change in neighborhood characteristics and change in coronary artery calcium: a longitudinal investigation in the MESA (Multi-Ethnic Study of Atherosclerosis) Cohort. *Circulation*. 2016;134:504–513. doi: 10.1161/CIRCULATIONAHA.115.020534.
64. Müller-Riemenschneider F, Pereira G, Villanueva K, Christian H, Knuiman M, Giles-Corti B, Bull FC. Neighborhood walkability and cardiometabolic risk factors in Australian adults: an observational study. *BMC Public Health*. 2013;13:755. doi: 10.1186/1471-2458-13-755.
65. Van Dyck D, Cerin E, Cardon G, Deforche B, Sallis JF, Owen N, de Bourdeaudhuij I. Physical activity as a mediator of the associations between neighborhood walkability and adiposity in Belgian adults. *Health Place*. 2010;16:952–960. doi: 10.1016/j.healthplace.2010.05.011.
66. Sampson RJ, Sharkey P. Neighborhood selection and the social reproduction of concentrated racial inequality. *Demography*. 2008;45:1–29.
67. Glass TA, Bilal U. Are neighborhoods causal? Complications arising from the 'stickiness' of ZNA. *Soc Sci Med*. 2016;166:244–253. doi: 10.1016/j.socscimed.2016.01.001.
68. Rhone A, Ploeg MV, Dicken C, Williams R, Breneman V. Low-income and low-supermarket-access census tracts, 2010–2015. EIB-165. Washington, DC: US Department of Agriculture, Economic Research Service; January 2017.
69. Sumner JA, Khodnava Y, Muntner P, Redmond N, Lewis MW, Davidson KW, Edmondson D, Richman J, Safford MM. Effects of concurrent depressive symptoms and perceived stress on cardiovascular risk in low- and high-income participants: findings from the Reasons for Geographical and Racial Differences in Stroke (REGARDS) Study. *J Am Heart Assoc*. 2016;5:1–14.
70. Lazzarino AI, Hamer M, Stamatakis E, Steptoe A. Low socioeconomic status and psychological distress as synergistic predictors of mortality from stroke and coronary heart disease. *Psychosom Med*. 2013;75:311–316. doi: 10.1097/PSY.0b013e3182898e6d.
71. Heslop P, Smith GD, Carroll D, Macleod J, Hyland A, Hart C. Perceived stress and coronary heart disease risk factors: the contribution of socioeconomic position. *Br J Health Psychol*. 2001;6(pt 2):167–178. doi: 10.1348/135910701169133.
72. Ng DM, Jeffery RW. Relationships between perceived stress and health behaviors in a sample of working adults. *Health Psychol*. 2003;22:638–642. doi: 10.1037/0278-6133.22.6.638.
73. Ye S, Muntner P, Shimbo D, Judd SE, Richman J, Davidson KW, Safford MM. Behavioral mechanisms, elevated depressive symptoms, and the risk for myocardial infarction or death in individuals with coronary

- heart disease: the REGARDS (Reason for Geographic and Racial Differences in Stroke) study. *J Am Coll Cardiol*. 2013;61:622–630. doi: 10.1016/j.jacc.2012.09.058.
74. Shaw LJ, Merz CN, Bittner V, Kip K, Johnson BD, Reis SE, Kelsey SF, Olson M, Mankad S, Sharaf BL, Rogers WJ, Pohost GM, Sopko G, Pepine CJ; WISE Investigators. Importance of socioeconomic status as a predictor of cardiovascular outcome and costs of care in women with suspected myocardial ischemia. Results from the National Institutes of Health, National Heart, Lung and Blood Institute-sponsored Women's Ischemia Syndrome Evaluation (WISE). *J Womens Health (Larchmt)*. 2008;17:1081–1092. doi: 10.1089/jwh.2007.0596.
 75. Fabreau GE, Leung AA, Southern DA, Knudtson ML, McWilliams JM, Ayanian JZ, Ghali WA. Sex, socioeconomic status, access to cardiac catheterization, and outcomes for acute coronary syndromes in the context of universal healthcare coverage. *Circ Cardiovasc Qual Outcomes*. 2014;7:540–549. doi: 10.1161/CIRCOUTCOMES.114.001021.
 76. LeFevre ML; U.S. Preventive Services Task Force. Behavioral counseling to promote a healthful diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med*. 2014;161:587–593. doi: 10.7326/M14-1796.
 77. Lin JS, O'Connor E, Evans CV, Senger CA, Rowland MG, Groom HC. Behavioral counseling to promote a healthy lifestyle in persons with cardiovascular risk factors: a systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2014;161:568–578. doi: 10.7326/M14-0130.
 78. Brown J, Michie S, Geraghty AW, Yardley L, Gardner B, Shahab L, Stapleton JA, West R. Internet-based intervention for smoking cessation (StopAdvisor) in people with low and high socioeconomic status: a randomised controlled trial. *Lancet Respir Med*. 2014;2:997–1006. doi: 10.1016/S2213-2600(14)70195-X.
 79. Siren R, Eriksson JG, Peltonen M, Vanhanen H. Impact of health counselling on cardiovascular disease risk in middle aged men: influence of socioeconomic status. *PLoS One*. 2014;9:e88959. doi: 10.1371/journal.pone.0088959.
 80. Jiang L, Huang H, Johnson A, Dill EJ, Beals J, Manson SM, Roubideaux Y; Special Diabetes Program for Indians Diabetes Prevention Demonstration Project. Socioeconomic disparities in weight and behavioral outcomes among American Indian and Alaska Native participants of a translational lifestyle intervention project. *Diabetes Care*. 2015;38:2090–2099. doi: 10.2337/dc15-0394.
 81. Blair SN, Kampert JB, Kohl HW 3rd, Barlow CE, Macera CA, Paffenbarger RS Jr, Gibbons LW. Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. *JAMA*. 1996;276:205–210.
 82. Sesso HD, Paffenbarger RS Jr, Lee IM. Physical activity and coronary heart disease in men: the Harvard Alumni Health Study. *Circulation*. 2000;102:975–980.
 83. Borrell LN. The effects of smoking and physical inactivity on advancing mortality in U.S. adults. *Ann Epidemiol*. 2014;24:484–487. doi: 10.1016/j.annepidem.2014.02.016.
 84. Schiller JS, Lucas JW, Ward BW, Peregoy JA. Summary health statistics for U.S. adults: National Health Interview Survey, 2010. *Vital Health Stat*. 2012;1–207.
 85. Andrade AC, Peixoto SV, Friche AA, Goston JL, César CC, Xavier CC, Proietti FA, Diez Roux AV, Caiaffa WT. Social context of neighborhood and socioeconomic status on leisure-time physical activity in a Brazilian urban center: The BH Health Study. *Cad Saude Publica*. 2015;31(suppl 1):136–147. doi: 10.1590/0102-311X00069514.
 86. Gidlow C, Johnston LH, Crone D, Ellis N, James D. A systematic review of the relationship between socio-economic position and physical activity. *Health Ed J*. 2006;65:338–367.
 87. Schulz AJ, Israel BA, Mentz GB, Bernal C, Caver D, DeMajo R, Diaz G, Gamboa C, Gaines C, Hoston B, Opperman A, Reyes AG, Rowe Z, Sand SL, Woods S. Effectiveness of a walking group intervention to promote physical activity and cardiovascular health in predominantly non-Hispanic black and Hispanic urban neighborhoods: findings from the walk your heart to health intervention. *Health Educ Behav*. 2015;42:380–392. doi: 10.1177/1090198114560015.
 88. Havranek EP, Mujahid MS, Barr DA, Blair IV, Cohen MS, Cruz-Flores S, Davey-Smith G, Dennison-Himmelfarb CR, Lauer MS, Lockwood DW, Rosal M, Yancy CW; American Heart Association Council on Quality of Care and Outcomes Research, Council on Epidemiology and Prevention, Council on Cardiovascular and Stroke Nursing, Council on Lifestyle and Cardiometabolic Health, and Stroke Council. Social determinants of risk and outcomes for cardiovascular disease: a scientific statement from the American Heart Association. *Circulation*. 2015;132:873–898. doi: 10.1161/CIR.0000000000000228.
 89. Pearson TA, Palaniappan LP, Artinian NT, Carnethon MR, Criqui MH, Daniels SR, Fonarow GC, Fortmann SP, Franklin BA, Galloway JM, Goff DC Jr, Heath GW, Frank AT, Kris-Etherton PM, Labarthe DR, Murabito JM, Sacco RL, Sasson C, Turner MB; American Heart Association Council on Epidemiology and Prevention. American Heart Association Guide for Improving Cardiovascular Health at the Community Level, 2013 update: a scientific statement for public health practitioners, healthcare providers, and health policy makers. *Circulation*. 2013;127:1730–1753. doi: 10.1161/CIR.0b013e31828f8a94.
 90. Jeemon P, Narayanan G, Kondal D, Kahol K, Bharadwaj A, Purty A, Negi P, Ladhani S, Sanghvi J, Singh K, Kapoor D, Sobti N, Lall D, Manimunda S, Dwivedi S, Toteja G, Prabhakaran D; DISHA Study Investigators. Task shifting of frontline community health workers for cardiovascular risk reduction: design and rationale of a cluster randomised controlled trial (DISHA study) in India. *BMC Public Health*. 2016;16:264. doi: 10.1186/s12889-016-2891-6.
 91. Adeyemo A, Tayo BO, Luke A, Ogedegbe O, Durazo-Arvizu R, Cooper RS. The Nigerian antihypertensive adherence trial: a community-based randomized trial. *J Hypertens*. 2013;31:201–207. doi: 10.1097/HJH.0b013e32835b0842.
 92. Gaziano T, Abrahams-Gessel S, Surka S, Sy S, Pandya A, Denman CA, Mendoza C, Puaane T, Levitt NS. Cardiovascular disease screening by community health workers can be cost-effective in low-resource countries. *Health Aff (Millwood)*. 2015;34:1538–1545. doi: 10.1377/hlthaff.2015.0349.
 93. Gaziano TA, Abrahams-Gessel S, Denman CA, Montano CM, Khanam M, Puaane T, Levitt NS. An assessment of community health workers' ability to screen for cardiovascular disease risk with a simple, non-invasive risk assessment instrument in Bangladesh, Guatemala, Mexico, and South Africa: an observational study. *Lancet Glob Health*. 2015;3:e556–e563. doi: 10.1016/S2214-109X(15)00143-6.
 94. Shroufi A, Chowdhury R, Anchala R, Stevens S, Blanco P, Han T, Niessen L, Franco OH. Cost effective interventions for the prevention of cardiovascular disease in low and middle income countries: a systematic review. *BMC Public Health*. 2013;13:285. doi: 10.1186/1471-2458-13-285.
 95. Afshin A, Penalvo J, Del Gobbo L, Kashaf M, Michal R, Morrish K, Pearson-Stuttard J, Rehm C, Shangquan S, Smith JD, Mozaffarian D. CVD prevention through policy: a review of mass media, food/menu labeling, taxation/subsidies, built environment, school procurement, worksite wellness, and marketing standards to improve diet. *Curr Cardiol Rep*. 2015;17:98. doi: 10.1007/s11886-015-0658-9.
 96. Interactive Atlas of Heart Disease and Stroke. Atlanta, GA: Centers for Disease Control and Prevention, Division for Heart Disease and Stroke Prevention. <http://www.cdc.gov/dhdsdp/maps/atlas>. Accessed February 22, 2018.
 97. Bagheri N, Gilmour B, McRae I, Konings P, Dawda P, Del Fante P, van Weel C. Community cardiovascular disease risk from cross-sectional general practice clinical data: a spatial analysis. *Prev Chronic Dis*. 2015;12:E26. doi: 10.5888/pcd12.140379.
 98. Engdahl J, Holmén A, Rosenqvist M, Strömberg U. Uptake of atrial fibrillation screening aiming at stroke prevention: geo-mapping of target population and non-participation. *BMC Public Health*. 2013;13:715. doi: 10.1186/1471-2458-13-715.
 99. Sampson UK, Kaplan RM, Cooper RS, Diez Roux AV, Marks JS, Engelgau MM, Peprah E, Mishoe H, Boulware LE, Felix KL, Califf RM, Flack JM, Cooper LA, Gracia JN, Henderson JA, Davidson KW, Krishnan JA, Lewis TT, Sanchez E, Luban NL, Vaccarino V, Wong WF, Wright JT Jr, Meyers D, Ogedegbe OG, Presley-Cantrell L, Chambers DA, Belis D, Bennett GC, Boyington JE, Creazzo TL, de Jesus JM, Krishnamurti C, Lowden MR, Punturieri A, Shero ST, Young NS, Zou S, Mensah GA. Reducing health inequities in the U.S.: recommendations from the NHLBI's Health Inequities Think Tank Meeting. *J Am Coll Cardiol*. 2016;68:517–524. doi: 10.1016/j.jacc.2016.04.059.
 100. Bajekal M, Scholes S, O'Flaherty M, Raine R, Norman P, Capewell S. Unequal trends in coronary heart disease mortality by socioeconomic circumstances, England 1982–2006: an analytical study. *PLoS One*. 2013;8:e59608.
 101. Lotufo PA, Fernandes TG, Bando DH, Alencar AP, Benseñor IM. Income and heart disease mortality trends in Sao Paulo, Brazil, 1996 to 2010. *Int J Cardiol*. 2013;167:2820–2823. doi: 10.1016/j.ijcard.2012.07.006.
 102. Kanjilal S, Gregg EW, Cheng YJ, Zhang P, Nelson DE, Mensah G, Beckles GL. Socioeconomic status and trends in disparities in 4 major risk factors for cardiovascular disease among US adults, 1971–2002. *Arch Intern Med*. 2006;166:2348–2355. doi: 10.1001/archinte.166.21.2348.

103. O'Flaherty M, Bishop J, Redpath A, McLaughlin T, Murphy D, Chalmers J, Capewell S. Coronary heart disease mortality among young adults in Scotland in relation to social inequalities: time trend study. *BMJ*. 2009;339:b2613.
104. Sims M, Diez Roux AV, Boykin S, Sarpong D, Gebreab SY, Wyatt SB, Hickson D, Payton M, Ekunwe L, Taylor HA. The socioeconomic gradient of diabetes prevalence, awareness, treatment, and control among African Americans in the Jackson Heart Study. *Ann Epidemiol*. 2011;21:892–898. doi: 10.1016/j.annepidem.2011.05.006.
105. Woodward M, Brindle P, Tunstall-Pedoe H; SIGN group on risk estimation. Adding social deprivation and family history to cardiovascular risk assessment: the ASSIGN score from the Scottish Heart Health Extended Cohort (SHHEC). *Heart*. 2007;93:172–176. doi: 10.1136/hrt.2006.108167.
106. Hippisley-Cox J, Coupland C, Vinogradova Y, Robson J, Minhas R, Sheikh A, Brindle P. Predicting cardiovascular risk in England and Wales: prospective derivation and validation of QRISK2. *BMJ*. 2008;336:1475–1482. doi: 10.1136/bmj.39609.449676.25.
107. Franks P, Tancredi DJ, Winters P, Fiscella K. Including socioeconomic status in coronary heart disease risk estimation. *Ann Fam Med*. 2010;8:447–453. doi: 10.1370/afm.1167.
108. Chao TF, Liu CJ, Wang KL, Lin YJ, Chang SL, Lo LW, Hu YF, Tuan TC, Chen TJ, Lip GY, Chen SA. Using the CHA2DS2-VASc score for refining stroke risk stratification in 'low-risk' Asian patients with atrial fibrillation. *J Am Coll Cardiol*. 2014;64:1658–1665. doi: 10.1016/j.jacc.2014.06.1203.
109. Chao TF, Liu CJ, Wang KL, Lin YJ, Chang SL, Lo LW, Hu YF, Tuan TC, Chen TJ, Lip GY, Chen SA. Should atrial fibrillation patients with 1 additional risk factor of the CHA2DS2-VASc score (beyond sex) receive oral anticoagulation? *J Am Coll Cardiol*. 2015;65:635–642. doi: 10.1016/j.jacc.2014.11.046.
110. Yusuf S, Wood D, Ralston J, Reddy KS. The World Heart Federation's vision for worldwide cardiovascular disease prevention. *Lancet*. 2015;386:399–402. doi: 10.1016/S0140-6736(15)60265-3.
111. Smith SC Jr, Chen D, Collins A, Harold JG, Jessup M, Josephson S, Logstrup S, Jur C, Sacco RL, Vardas PE, Wood DA, Zoghbi WA. Moving from political declaration to action on reducing the global burden of cardiovascular diseases: a statement from the global cardiovascular disease taskforce. *Circulation*. 2013;128:2546–2548. doi: 10.1161/CIR.0b013e3182a93504.
112. Vanassche T, Connolly SJ, Eikelboom JW, Healey JS, Lauw MN, Masiero S, Wang J, Yusuf S. Abstract 18936: Can biomarkers improve risk stratification of atrial fibrillation patients? Analysis of 3578 aspirin-treated patients in ACTIVE and AVERROES. *Circulation*. 2014;130:A18936.
113. Williams DR, Purdie-Vaughns, V. Social and behavioral interventions to improve health and reduce disparities in health. <https://www.ahrq.gov/professionals/education/curriculum-tools/population-health/williams.html>. Published 2015. Accessed January 30, 2018.
114. Tobias M. Social rank: a risk factor whose time has come? *Lancet*. 2017;389:1172–1174. doi: 10.1016/S0140-6736(17)30191-5.