

dfghjhgfrtyu

by Oihgfdghj Dfghjk

Submission date: 27-Apr-2020 11:42AM (UTC+0300)

Submission ID: 1308958527

File name: synopsis.edited.edited.edited.docx (18.12K)

Word count: 976

Character count: 5469

Running head: ¹DIABETES MELLITUS IS AN INDEPENDENT PROGNOSTIC FACTOR FOR TAVI 1

¹Diabetes Mellitus Is an Independent Prognostic Factor for TAVI

Student's Name

Instructor's Name

Date

¹ Diabetes Mellitus Is an Independent Prognostic Factor for TAVI

Diabetes mellitus has been increasing in the world and causing increased mortality rates. Diabetes mellitus is a metabolic disease which leads to an increase in blood sugar. A healthy body works through the insulin hormone moving the sugar from the blood to the cells for it to be used as energy or stored. However, while having diabetes mellitus disease, the body can neither make the required insulin nor can it efficiently use the already made insulin. Therefore, when the high blood sugar generated by diabetes disease is not treated, it may lead to damage to body organs such as the eyes, nerves and kidneys. Due to the increased negative roles that diabetes mellitus has played during the surgical process; the article has focused on patients who have undergone transcatheter aortic valve implantation.

Similarly, the article has researched on the Pubmed database to find the effects that diabetes mellitus has had both long term, short term and mid-term diagnosis to patients who have experienced ⁶ transcatheter aortic valve replacement. Transcatheter aortic valve replacement is a nominal aggressive clinical practice for repairing the valve without removing it. Diabetes mellitus is considered as the leading cause in rise of cardiovascular deaths as well as various complications like coronary heart disorders which lead to an intensification in healthcare cost reduction in life expectancy. Aortic stenosis is regarded as the most known ⁴ valvular heart disease in developed countries and is expected to increase due to the increase in the aging population. Nonetheless, surgical aortic valve replacement is a common standard therapeutic choice; nevertheless, old patients are regarded as vulnerable to surgical and hence barred from using the therapy.

Since the study is performed by following the required standards provided by the systematic reviews and meta-analysis reporting, every study used is required to certify the

eligibility of participants involved in the study which are needed to be aortic stenosis (AS), an intervention which requires only transcatheter aortic valve implantation (TAVI). Similarly, the outcome is needed to provide short-term, long-term and mid-term persistence rate for patients who endure TAVI. The article used a search strategy of finding previous research from the PubMed electronic database, where articles from 1st March 2017 to 15th March 2017 were searched using the keyword TAVI and Diabetes.

The quality of the information is vital to provide relevant and useful information. As a result, the Newcastle-Ottawa Scale is used as a methodology to evaluate the quality of non-randomized studies. The analysis is done by evaluating the diabetes mellitus mortality rate within thirty days, where a ninety-five percent confidence level of odds ratio is used to predict the qualitative synthesis. Similarly, the hazard ratio of ninety-five percent CI is used as the conclusion of the prediction for both long-term and overall mid-term survivor and the hazard ratio is the only suitable statistic that is well-matched time to event value and expurgating. The results are then synthesized using I^2 statistics and Cochrane's Q-test. At the same time, the heterogeneity level is being monitored such that when it reaches a point of fine heterogeneity with p being more significant than 0.1 and I^2 being less than fifty percent, then the **standard** **fixed-effect model test** can be utilized. Else the random **fixed-effect model test** could be of help.

Despite identifying four hundred and sixty-seven citations from PubMed database and five studies from other sources, only twenty studies are used in the study. To evaluate the outcome of diabetes mellitus on the diagnosis of TAVI, the data concerning both medium and long-term OS, is further divided into several subgroups based on the background of an individual, clinic period and numerical investigation. Publications for long term OS indicates that diabetes mellitus could be extrapolative for both poor long term and medium term for people

who have experienced TAVI. However, from the subgroups, only ten publications offer results from multivariable scrutiny, and it designated that diabetes mellitus is related to both lower medium to long term OS. Moreover, from the fourteen publications three are based on Asian individuals while eleven are based on western individuals. The projections indicate a suggestively ¹ lower medium to long term OS in western individuals having diabetes Mellitus. At the same time, there is no expressive association indicated by diabetes mellitus and prognosis of TAVI in Asian individuals.

According to the study, most publications indicate that it is more favorable for medium to long term OS in individuals free from diabetes mellitus to be performed TAVI than to those with diabetes mellitus. Similarly, diabetes mellitus is regarded as the main forecaster of medium-term existence for TAVI process. Nevertheless, diabetic Mellitus is also found to be related to ³ the long-term death of TAVI. However, based on the research, there is no strong effect of ³ diabetes mellitus indicated on thirty-day mortality, therefore, the projecting rate for diabetes mellitus for ¹ thirty-day mortality can be less than medium to long-term OS. Such an option is considered for the event to occur where individuals with diabetes mellitus could attain recommendable glycaemic control and thus be provided with a more antagonistic action relating to diabetes mellitus disorders.

Several limitations are encountered during the study; first, the use of randomized control trials is used in meta-analysis due to the topic complexity since it was not probable to group individuals based on their diabetic position. Secondly, although there is no journal bias, journals that report positive outcomes are always more expected to be disseminated than those reporting negative effects.

Reference

- ² Lv, W., Li, S., Zhao, Z., Liao, Y., Li, Y., Chen, M., & Feng, Y. (2018). Diabetes mellitus is an independent prognostic factor for mid-term and long-term survival following transcatheter aortic valve implantation: a systematic review and meta-analysis. *Interactive cardiovascular and thoracic surgery*, 27(2), 159-168. Retrieved from ³ <https://academic.oup.com/icvts/article/27/2/159/4924811>

ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1

academic.oup.com

Internet Source

6%

2

academic.oup.com

Internet Source

4%

3

Wenyu Lv, Shuangjiang Li, Zhengang Zhao, Yanbiao Liao, Yijian Li, Mao Chen, Yuan Feng. "Diabetes mellitus is an independent prognostic factor for mid-term and long-term survival following transcatheter aortic valve implantation: a systematic review and meta-analysis", Interactive CardioVascular and Thoracic Surgery, 2018

Publication

3%

4

worldwidescience.org

Internet Source

1%

5

pesquisa.bvsalud.org

Internet Source

1%

6

www.dovepress.com

Internet Source

1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off