

UC Riverside

UC Riverside Previously Published Works

Title

Upregulating the positive affect system in anxiety and depression: Outcomes of a positive activity intervention

Permalink

<https://escholarship.org/uc/item/06v1w2v1>

Journal

DEPRESSION AND ANXIETY, 34(3)

ISSN

1091-4269

Authors

Taylor, CT
Lyubomirsky, S
Stein, MB

Publication Date

2017-03-01

DOI

10.1002/da.22593

Peer reviewed

Running Head: PAI RCT FOR ANXIETY AND DEPRESSION – TAYLOR

Upregulating the Positive Affect System in Anxiety and Depression:

Outcomes of a Positive Activity Intervention

Charles T. Taylor^{a,*}

Sonja Lyubomirsky^b

Murray B. Stein^a

^a University of California, San Diego

^b University of California, Riverside

in press, *Depression and Anxiety*

* Corresponding author:

Charles T. Taylor, Ph.D.
Department of Psychiatry
University of California, San Diego
La Jolla, CA 92037
Phone: 858.534.9446
FAX: 858.534.9450
Email: c1taylor@ucsd.edu

Funding: This research was supported by a grant awarded to Charles T. Taylor from the University of California, San Diego, National Institute of Health Clinical and Translational Science Awards Program Grant UL1TR001442.

Abstract

Background: Research suggests that the positive affect system may be an important yet underexplored treatment target in anxiety and depression. Existing interventions primarily target the negative affect system, yielding modest effects on measures of positive emotions and associated outcomes (e.g., psychological well-being). The objective of the present pilot study was to evaluate the efficacy of a new transdiagnostic positive activity intervention (PAI) for anxiety and depression.

Method: Twenty-nine treatment-seeking individuals presenting with clinically impairing symptoms of anxiety and/or depression were randomly allocated to a 10-session protocol comprised of PAIs previously shown in non-clinical samples to improve positive thinking, emotions, and behaviors (e.g., gratitude, acts of kindness, optimism; $n=16$) or a waitlist condition ($n=13$). Participants were assessed at pre- and post-treatment, as well as 3- and 6-month follow-up, on measures of positive and negative affect, symptoms, and psychological well-being.

ClinicalTrials.gov Identifier:NCT02330627

Results: The PAI group displayed significantly larger improvements in positive affect and psychological well-being from pre- to post-treatment compared to waitlist. Post-treatment and follow-up scores in the PAI group were comparable to general population norms. The PAI regimen also resulted in significantly larger reductions in negative affect, as well as anxiety and depression symptoms, compared to waitlist. Improvements across all outcomes were large in magnitude and maintained over a 6-month follow-up period.

Conclusions: Targeting the positive affect system through a multi-component PAI regimen may be beneficial for generating improvements in positive emotions and well-being, as well as reducing negative affect and symptoms, in individuals with clinically impairing anxiety or

depression.

Keywords: Anxiety; depression; transdiagnostic; positive activities; positive intervention; positive affect; well-being; randomized controlled trial

Upregulating the Positive Affect System in Anxiety and Depression:

Outcomes of a Positive Activity Intervention

Anxiety and depressive disorders are the most common mental health conditions¹ and represent a major public health concern worldwide^{2, 3}. These conditions frequently co-occur⁴, significantly impair functioning, and diminish quality of life and well-being^{5, 6}. Recent efforts to integrate advances in neuroscience with clinical psychiatry suggest that anxiety and depression can be conceptualized along partially distinct biobehavioral dimensions of positive and negative affect domains^{7, 8} [Footnote 1]. The negative affect system regulates responses to cues that signal potential danger or loss, and is characterized by negative emotions (e.g., fear, sadness), cognitions (e.g., rumination), and inhibitory/avoidance behaviors. In contrast, the positive affect system guides people toward situations with reward potential, and is characterized by positive emotions (e.g., joy, excitement, happiness), cognitions (e.g., attentional bias for reward-relevant stimuli), and approach behaviors (e.g., curiosity, social initiation) that together facilitate the acquisition of psychosocial resources that promote overall health and well-being^{9, 10}. Features of the negative affect system are centrally positioned within prevailing diagnostic classification systems¹¹ and conceptual models of anxiety and depression^{12, 13}, and thus serve as primary targets of existing psychosocial intervention approaches¹⁴⁻¹⁸. However, accumulating research suggests that the positive affect system may serve as an important yet underexplored target in facilitating recovery from anxiety and depression^{9, 19-21}.

Positive emotions serve a number of functions that both mitigate the adverse effects of negative emotions, the defining features of anxiety and depressive disorders, as well as garner positive outcomes that promote resilience and psychological well-being. For example, positive

emotions down-regulate the physiological and psychological effects of negative emotions²²⁻²⁵ – theorized to occur in part through counteracting the narrow, inflexible, and negatively biased patterns of cognition and behavior (e.g., avoidance) that perpetuate negative mood states²⁶⁻³⁰ (see⁹ for a review). The positive affect system also fosters approach-oriented behaviors, such as exploration and social initiation, that increase exposure to potentially rewarding outcomes,³¹ thereby facilitating the acquisition of social, physical, and intellectual resources^{9, 10, 32} that promote resilience during stress and overall well-being³³. The unique link between positive emotions and subjective well-being above and beyond negative affect and psychopathology symptoms^{34, 35} suggests that interventions targeting the positive emotion system may fill a particularly important gap left by extant treatments.

Evidence across multiple units of analysis reveals that depression¹⁹, and some forms of anxiety (e.g., social anxiety^{36, 37}; posttraumatic stress disorder³⁸) are associated with aberrant functioning of the positive emotion system, including low positive affect³⁹⁻⁴¹, diminished approach motivation and behavior⁴²⁻⁴⁶, biased processing of reward-related cues⁴⁷⁻⁵⁰ and reduced behavioral⁵¹ and neural reactivity to rewards⁵²⁻⁵⁵ (for reviews see^{8, 56}). Existing interventions for anxiety and depression emphasize negative affect reduction as the central treatment goal. Given that positive and negative affect arise from partially distinct biobehavioral systems^{57, 58}, decreases in negative affect and symptoms may not result in concomitant increases in positive affect and related outcomes. Consistent with this proposal, prevailing intervention approaches for anxiety and depression display modest effects on the positive affect system and associated outcomes (e.g., psychological well-being)^{6, 59-62}, and in some cases, have failed to show evidence of changes following treatment, despite significant reductions in anxiety and depression⁶³. For example, a large randomized controlled trial comparing cognitive therapy, pharmacotherapy, and

their combination for depression revealed that participants displayed post-treatment levels of positive affect that were significantly below community norms, despite returning to normative levels of negative affect⁶⁴. Thus, the limited evidence that exists suggests that current intervention approaches for anxiety and depression may not be sufficiently targeting the biobehavioral processes that are important for building positive emotions and well-being.

The central aim of the current pilot study was to test the efficacy of a psychosocial treatment protocol designed to upregulate the positive affect system in a sample of individuals seeking treatment for anxiety or depression. The intervention is grounded in research demonstrating that people can increase their levels of positive thinking, emotions, and behavior through engaging in simple, intentional, and repeated activities^{65, 66}, for example, performing acts of kindness, expressing gratitude, and practicing optimism (see Table 1). Studies supporting the efficacy of positive activity interventions (PAIs) have been primarily conducted in unselected (non-clinical) community samples and examined single positive activities in isolation (for a meta-analysis, see^{66, 67}). Although some evidence suggests that integrated, multi-session PAIs may be beneficial for individuals with mild-to-moderate depression⁶⁸ or other health-related conditions (e.g., cardiovascular disease⁶⁹, suicide⁷⁰, and HIV⁷¹), to our knowledge, no studies have examined PAIs in psychiatric treatment-seeking samples. That was the central goal of the current study.

Method

Participants

The sample comprised $N=29$ individuals seeking treatment for anxiety and/or depression. Participants were recruited through clinical referrals as well as posted announcements in community and online settings (e.g., ResearchMatch.org). Participants were required to be

between the ages of 18 to 55 and present with clinically elevated symptoms of anxiety or depression, defined by a score of 8 or higher on the Overall Anxiety Severity and Impairment Scale (OASIS⁷²) or a score of 10 or higher on the Patient Health Questionnaire-9 (PHQ-9⁷³), respectively. Exclusion criteria were used to ensure that participants could safely complete the procedures and to minimize confounding interpretation of our findings^[Footnote 2]: (1) pharmacological treatments that could affect brain functioning (e.g., anxiolytics, antidepressants; past 6 weeks); (2) concurrent psychotherapy, or empirically supported treatments for anxiety or depression (e.g., cognitive behavioral therapy; past 6 weeks); (3) current active suicidal ideation; (4) history of major neurological disorder or moderate to severe traumatic brain injury; (5) moderate alcohol or marijuana use disorder (past year); mild substance use disorder (all other drugs; past year); (6) bipolar I or psychotic disorders; and (7) characteristics that compromise MRI safety (e.g., metal in body).

Diagnostic assessment was based on a structured diagnostic interview for DSM-5, Mini International Neuropsychiatric Interview (MINI Version 7.0.0.0). Participant enrollment statistics and progress through the study are summarized in Figure 1. Participants were enrolled in the study between June 2014 and March 2015. Of the 29 participants who were randomized to the PAI ($n=16$) or waitlist group ($n=13$), one participant in the PAI group discontinued treatment following session 7, reportedly due to increased commitments at work, and one participant in the waitlist condition initiated treatment following the pre-assessment and was excluded from the analyses^[Footnote 3]. Thus, 28 participants ($n=16$ in the PAI group and $n=12$ in the waitlist group) were included in the intent-to-treat analysis. The demographic and clinical composition reflected a diverse, community-based treatment-seeking sample (see Table 2).

Measures

Participants completed a battery of reliable and valid self-report measures at pre-, post-, and follow-up assessment points (3 and 6 months following the post-assessment).

Positive and Negative Emotions. Participants completed the 20-item Positive and Negative Affect Schedule - Trait (PANAS)⁷⁴ to assess activated forms of positive and negative affect; and the 20-item Modified Differential Emotions Scale (mDES)²⁴ to assess a broader array of discrete positive (e.g., joy, love, awe) and negative emotions (e.g., guilt, anger, fear).

Participants responded according to how they felt *during the past week*.

Psychological Well-Being. Participants completed the Quality of Life, Enjoyment, and Satisfaction Questionnaire – Short Form (Q-LES-Q-SF)⁷⁵ to measure perceived overall enjoyment and satisfaction across numerous life domains (e.g., work, health, relationships)⁷⁶; and the Satisfaction With Life Scale (SWLS)⁷⁷, a well-established measure of global judgments of satisfaction with one's life^{78, 79}.

Symptoms. Anxiety symptoms were assessed using the Overall Anxiety Severity and Impairment Scale (OASIS^{72, 80}), a 5-item scale that measures the frequency and severity of anxiety symptoms, as well as level of avoidance, work/school/home interference, and social interference associated with anxiety during the past 2 weeks; and the State Trait Anxiety Inventory-Trait (STAI-T⁸¹), a well-established measure of general anxiety. Depressed mood during the past 2 weeks was assessed using the Patient Health Questionnaire-9 (PHQ-9^{73, 82}) and the Beck Depression Inventory-II (BDI-II^{83, 84}).

Treatment Credibility/Expectancy. Following presentation of the treatment rationale, participants in the PAI group completed the Credibility and Expectancy Questionnaire⁸⁵, which asks about the logic of the intervention and its perceived likelihood of helping the participant and other people with anxiety or depression.

Procedure

Potential participants were given information about the study and provided informed written consent prior to completing the screening procedures, which comprised a MINI diagnostic interview and self-report assessments of anxiety and depression symptoms. Participants who met inclusion criteria and agreed to participate in the study were invited to complete a baseline evaluation session comprising self-report and behavioral assessments followed by a separate functional magnetic resonance imaging (fMRI) session, results of which will be reported separately. Following the MRI session, participants were randomly assigned to either the PAI group—that is, immediate treatment, or waitlist. Condition assignment was determined using a random number generator and revealed at the conclusion of the MRI session. Participants assigned to the PAI group completed 10 one-hour weekly sessions of the PAI protocol described below. Following the final treatment session (or approximately 10 weeks after the baseline MRI scan for the waitlist group), participants completed post-assessment sessions, which were identical to the pre-assessments. To establish the duration of treatment effects, participants in the PAI group completed self-report assessments 3- and 6-months following the post-assessment session. Waitlist participants were offered the PAI protocol following the post-assessment; however, their treatment data were not included in the analyses. Participants received monetary compensation for their participation in the assessment sessions. The procedures were approved by the University's Human Research Protections Program. ClinicalTrials.gov Identifier: NCT02330627

Treatment

Positive Activity Intervention (PAI). The PAI comprised 10 one-hour sessions of individual therapist-delivered treatment (plus a 30-minute introductory module at the start of the

first session to acquaint the therapist and patient, including a brief review of symptoms, past treatment experiences, and participant expectations and goals for treatment). A 71-page manual⁸⁶ that described the regimen in detail was developed based on prior literature on PAIs^{21, 69-71, 87, 88} and emotion science findings regarding the function of positive thoughts, emotions, and behaviors^{9, 20, 89-91}. Handouts accompanied each module, which included instructions for completing a given activity and text-fillable fields to allow participants to plan the activity and set goals, generate responses to the activity, and monitor their progress and observations.

Treatment began with education about the function of positive and negative emotions, and how anxiety and depression can disrupt positive experiences²⁰. Emphasis was placed on generating upward spirals of positive thinking, emotions, and behaviors as a means to overcoming anxiety and depression. The core treatment exercises included specific PAIs designed to increase positive thinking, emotions, and/or behavior. See Table 1 for details. The final module involved developing a personalized positivity activity plan wherein participants prepared for continued engagement in activities, building on gains made in treatment, and identifying strategies to minimize relapse. The structure of each session followed traditional behavioral treatment regimens—namely, review completion of the prior week’s exercises, including self-monitoring of emotions and exercise completion; troubleshoot issues that arose during exercise completion; introduce material about a new PAI; and identify exercises to implement for the upcoming week.

Waitlist (WL). Waitlist participants completed the pre- and post-assessments at a 10-week interval. Treatment was offered to these individuals after the post-assessment.

Therapists. Therapists were one doctoral-level and one master’s level clinician, each with over 10 years of experience treating individuals with anxiety or depression. Both therapists

contributed to the development of the treatment protocol. Therapists met weekly to review ongoing cases to ensure that treatment material outlined in the manual was being appropriately covered and to discuss issues that arose in treatment. Treatment adherence was closely monitored during weekly supervision; however, it was not formally evaluated.

Statistical Analyses

Our primary outcome was positive emotions (composite of PANAS-PA and mDES-Positive Emotion scores). Secondary outcomes were: (1) negative emotions (composite of PANAS-NA and mDES-Negative Emotion scores); (2) psychological well-being (composite of Q-LES-Q-SF and SWLS scores); (3) anxiety (composite of OASIS and STAI-T scores); and (4) depression (composite of PHQ-9 and BDI-II scores). Analyses were conducted on an intent-to-treat (ITT) basis (PAI, $n=16$; WL, $n=12$). For the one participant who discontinued treatment following session 7, we used measures completed at a mid-treatment (i.e., the last available assessment point) for post-treatment data ^[Footnote 4].

Analysis of covariance (ANCOVA) was used to test group differences at post-treatment controlling for participants' pre-treatment scores for the measure of interest^{92, 93}. We tested and confirmed that all assumptions underlying ANCOVA were met (e.g., independence of the covariate [pre-treatment scores] and treatment group; homogeneity of regression slopes such that the covariate and treatment group do not interact in predicting the outcome). Following prior studies⁹⁴⁻⁹⁶, conceptually related measures were combined using Rosenthal and Rosnow's⁹⁷ procedure to create a composite index for each outcome domain. This approach arguably creates a more robust outcome index, and constrains type I error rate inflation. Participants' scores on each scale were first standardized ($M = 0$, $SD = 1$) across assessment sessions by converting to Z scores. The composite index at each assessment point was the mean of the Z scores for that

occasion. The magnitude of treatment response was established by calculating (a) within-group effect sizes = $([\text{post-assessment mean} - \text{pre-assessment mean}] / [\text{pre-assessment standard deviation} + \text{post-treatment standard deviation}] / 2)$ (referred to as Cohen's d_{av} ; see ⁹⁸); and (b) between-group controlled effect sizes = $(\text{post-assessment PAI covariance adjusted mean} - \text{post-assessment WL covariance adjusted mean}) / \text{pooled standard deviation}$. All analyses were conducted using SPSS version 18.0.

Results

Preliminary Analyses

Table 2 presents descriptive data for demographic and clinical characteristics for each group at baseline. Groups did not differ on gender, age, years of education, race, ethnicity, or past psychotropic medication use (all $ps > .10$). However, participants in the PAI group were significantly more likely to have reported prior psychosocial treatment use compared to participants in the waitlist group, Fisher's exact test significant = .044 ^[Footnote 5]. Credibility and Expectancy Questionnaire ratings obtained from participants in the PAI group following the first treatment session revealed high treatment rationale credibility ($M = 7.6$, $SD = .96$) and expectancy ($M = 6.8$, $SD = 2.0$; range of possible scores = 1-9).

Main Treatment Effects

Table 3 presents the means, standard deviations, and results of the ANCOVAs for the main outcomes. Results of the ANCOVA for our primary outcome revealed that individuals in the PAI group demonstrated significantly greater positive emotions at post-treatment compared to participants in the WL group ^[Footnote 6]. ANCOVA results for our secondary outcomes revealed that the PAI group reported experiencing significantly fewer negative emotions and symptoms of anxiety and depression, as well as significantly greater psychological well-being, at post-

treatment relative to participants in the WL group. The magnitude of both within- and between-group treatment effects was large for the PAI group. See Table 4.

Maintenance of Treatment Gains

Table 5 presents the means, standard deviations, and results of the repeated measures ANOVAs (Time: pre, post, 3- and 6-month follow-up) conducted on the main outcomes in the PAI group ^[Footnote 7]. All treatment completers ($n=15$) finished at least one follow-up assessment [$n=14$ at 3- and 6-month follow-up sessions]. Missing data at a given follow-up assessment point were substituted using data from a participant's last available assessment point (i.e., last observation carried forward). Results of the repeated measures ANOVAs conducted on the composite outcome indices all revealed significant main effects of Time. Follow-up contrasts using the Sidák adjustment for multiple comparisons indicated that the PAI group displayed significant changes on all outcome measures from pre- to post-assessment (all $ps < .05$), and from pre-assessment to each of the follow-up assessment points (all $ps < .05$). Post-treatment and follow-up scores did not significantly differ (all $ps > .05$), which indicated that initial gains were maintained up to 6-months following the end of treatment.

Discussion

The positive affect system is increasingly recognized as a potentially valuable treatment target for psychiatric conditions traditionally defined by heightened negative emotions, including anxiety and depression ¹⁹⁻²¹. We developed a multi-component protocol comprised of positive activity interventions (PAIs) designed to upregulate the positive emotion system, and took the first step in evaluating the efficacy of this regimen in individuals seeking treatment for anxiety or depression. The high treatment credibility and expectancy ratings as well as high completion rate suggests that the treatment regimen was well-received by the current sample, many of whom had

received prior psychosocial and/or pharmacological treatment. The PAI protocol resulted in significantly greater increases in positive emotions and psychological well-being compared to a no intervention control group. Interestingly, targeting the positive affect system generalized to negative affect-related outcomes, including reductions in negative emotions, anxiety, and depression. Treatment effects were large in magnitude and persisted up to 6-months following termination of acute treatment. Thus, the current preliminary findings underscore the potential value of directly targeting the positive affect system in treatment¹⁹⁻²¹, and add to a nascent empirical literature^{99, 100} suggesting that PAIs may be beneficial for individuals with clinically impairing symptoms of anxiety or depression.

The current sample was characterized by low levels of positive affect, and participants were, on average, dissatisfied with their lives upon entering treatment, scoring more than one standard deviation below normative general population means on both outcomes^{101, 102 103}. Following treatment, and persisting through the follow-up period, the PAI group scored near community normative means, suggesting that the PAI protocol restored positive emotional functioning to normative levels. Those findings are notable when considering that existing empirically supported interventions that primarily target the negative affect system display only modest effects in increasing positive emotions and well-being, even when resulting in significant and sometimes large reductions in negative affect^{63, 64}. Not all participants in the current study, however, achieved average or higher levels of life satisfaction following treatment, suggesting that additional PAI sessions or alternative treatment may be needed for some people. Nevertheless, in light of existing treatment outcome data, the current findings are promising and suggest that the PAI regimen is worthy of further empirical scrutiny.

Negative affect and symptoms of depression and anxiety were not direct treatment targets. However, the PAI protocol resulted in large reductions in negative affect and symptoms, changes that were comparable to those observed with prevailing empirically established interventions^{14, 15}. Those findings are consistent with prior studies demonstrating that positive emotions downregulate the adverse effects of negative emotions^{22, 23}, thinking (e.g., rumination³⁰), and help people cope during times of stress²⁴. It should be noted, however, that we administered broad measures of anxiety and depression, rather than assessing disorder-specific symptoms (e.g., panic, worry), and we did not conduct diagnostic assessments following treatment. Future research is needed to evaluate the effect of the PAI regimen on specific psychiatric conditions and symptom clusters, as well as to examine whether differential treatment response exists across disorders.

The current findings have clinical implications for emerging dimensional classification systems wherein the positive affect system is hypothesized to play a role across various forms of psychopathology, including anxiety and depression^{7, 8}. Consistent with these conceptual models, our goal was to develop a transdiagnostic protocol that could readily be applied to a range of psychiatric conditions, including subsyndromal cases that fail to meet diagnostic thresholds but may nevertheless experience marked functional impairment. Although the background material of the current intervention was tailored to anxiety and depression, the activities themselves were agnostic about psychopathology or specific symptom domains. Thus, the PAI protocol could, with slight modifications, be applied to other forms of psychopathology characterized by heightened negative affect and/or blunted positive affect. Moreover, as noted elsewhere¹⁰⁴⁻¹⁰⁶, transdiagnostic protocols have several advantages, including parsimony, reduced time and effort

to train providers, and facilitating clinicians' ability to treat comorbid clinical presentations typically seen in community practice.

The current findings should be interpreted in the context of several caveats. First, the efficacy of the PAI protocol was evaluated in a small treatment-seeking sample in comparison to an assessment only condition. Small sample sizes are prone to sampling biases such that the chosen sample may not be representative of the target population, and may produce outcomes that over- or underestimate the true treatment effect. Future studies are needed in larger samples using more rigorous comparison groups to account for common therapeutic effects (e.g., therapist attention, patient expectations of improvement), as well as to evaluate potentially unique mechanisms of change (e.g., positive emotions) compared to established interventions. Second, although therapists followed a structured manual and were monitored closely during weekly supervision, treatment fidelity and adherence were not formally assessed. Third, treatment outcomes relied on participant report. Although several of the central outcomes were, by definition, subjective in nature (e.g., well-being), future research is needed using clinician-administered and other objective (e.g., behavioral) measures. Moreover, indices of clinically significant change¹⁰⁷ would provide a more standardized benchmark for establishing reliable individual participant treatment response. Research is needed to establish test-retest reliability estimates of the primary (positive emotion) outcome measures in the target population and over a timeframe comparable to the pre-post interval used in the current (and many other treatment outcome) studies. Fourth, outcomes were only examined at pre- and post-treatment, and during follow-up. Future studies should include multiple repeated assessments of the outcome variables throughout treatment and use longitudinal statistical models (e.g., multilevel modeling), which are the preferred statistical approach for clinical trials. Such models honor the ITT principle and

are favored over other methods used to handle attrition (e.g., last observation carried forward) that may over- or underestimate treatment effects. Larger sample sizes (cf. the current study) are needed to appropriately conduct multilevel modeling using longitudinal data¹⁰⁸.

It is notable that the waitlist group displayed medium-sized changes on the positive and negative emotion outcome measures, but minimal changes on symptoms and well-being. Those outcomes may reflect ordinary fluctuations in emotions (cf. symptoms or well-being), natural improvement in treatment-seeking individuals (e.g., treatment expectancy effects¹⁰⁹), or sampling bias. Evaluating the PAI protocol in larger samples compared against active control conditions would help resolve these issues. Finally, concerns have been raised that PAIs may have adverse effects in some clinical disorders (e.g., depression) because focusing on positivity may invalidate one's suffering and distress (the Pollyanna problem¹⁹). Although we did not formally assess participant reactions to the individual positive activities, there was no evidence that participants deteriorated as a result of the intervention as a whole. The introductory psychoeducation session clarified that the aim of the intervention was *not* to feel positive emotions all of the time, nor to deny the existence of negative emotions or life experiences. Moreover, participants had freedom in choosing the activities that they ultimately incorporated into their daily lives. Nevertheless, the issue of person-activity fit^{87, 110} and treatment personalization¹¹¹ is an important one in need of future study.

Conclusions

The benefits of positive emotions^{9, 10} and their potential clinical utility for psychopathology¹⁹⁻²¹ are increasingly well-documented. Results of the current pilot study provided initial support for the efficacy of a multi-component treatment protocol comprised of PAIs in increasing positive emotions and psychological well-being, as well as decreasing

negative affect, anxiety, and depression in a treatment-seeking sample. Those findings support the potential value of explicitly targeting the positive emotion system in disorders classically defined and conceptualized according to heightened negative emotions. Future research in larger samples using more rigorous control conditions and moving beyond self-report measures is now needed.

Footnotes

¹Although we refer to positive and negative affect systems throughout the paper for consistency, we are speaking to the broader empirical literature on bivariate theories of human motivation and behavior^{7, 31, 74, 112-116}

²Participants completed a functional magnetic resonance imaging (fMRI) scan at pre- and post-assessment (data presented elsewhere). Hence, several of the exclusion criteria were implemented to ensure MRI safety and minimize confounding of the imaging findings.

³We conducted a sensitivity analysis in which the participant who initiated treatment during the waitlist period was included. Results of the ANCOVAs conducted on the main outcome composite indices revealed a nearly identical pattern of findings to those reported in the main text: The PAI group significantly differed from the WL group on all outcomes (all $ps < .05$); the only exception being that the treatment group effect on the positive emotion composite was marginally significant ($p = .065$). The between-group effect sizes (Cohen's d) on the post-treatment covariance adjusted scores ranged from 0.72 to 1.28, which is comparable to those reported in Table 4.

⁴We conducted a sensitivity analysis in which the participant who discontinued treatment prematurely was removed from the analysis. Results of the ANCOVAs conducted on the main outcome composite indices revealed the same pattern of findings to those reported in the main text: The PAI group significantly differed from the WL group on all outcomes (all $ps < .05$), and the between-group effect sizes (Cohen's d) on the post-treatment covariance adjusted scores were similar to those reported in Table 4 (range = 0.95 to 1.53).

⁵We conducted a sensitivity analysis for the main outcome variables in which history of prior psychosocial treatment was included as a covariate. Results revealed that psychosocial treatment history did not account for group differences observed across the main outcome indices.

⁶We also examined post-treatment (covariance adjusted) between-group effect size estimates of specific positive emotion items on the mDES to determine whether, for example, low activation positive emotions were more affected by the intervention. For 9 of 10 positive emotion items, results revealed medium to large between-group effects (Cohen's d range = 0.54 to 1.02), with lower activation positive emotion items showing large effects comparable to the full mDES positive emotion index (i.e., serene/content/peaceful, $d = 0.82$; grateful/appreciative/thankful, $d = 0.79$). The only exception to this pattern of findings was the item, amused/fun-loving/silly, which yielded a small between-group effect ($d = 0.17$).

⁷Given that the waitlist group did not complete follow-up assessments, composite scores used for the follow-up analyses were computed in the PAI group only across the four assessment sessions. Thus, mean composite index values differed for the PAI group for the between-group comparisons versus the follow-up analysis.

Acknowledgements

This research was supported by a grant awarded to Charles T. Taylor from the University of California, San Diego, Clinical and Translational Research Institute's National Institute of Health Clinical and Translational Science Awards Program Grant UL1TR001442.

We would like to thank the many individuals who helped make this research possible: Shadha Cissell for serving as a study therapist and her contributions to treatment manual development; Kristin Layous, Katie Nelson, Peter Ruberton, and Christina Armenta for providing valuable feedback on a draft version of the treatment manual content; Judith Moskowitz and Jeff Huffman for graciously sharing their positive affect intervention protocols developed for people adapting to chronic illness and patients hospitalized for suicidal ideation or behavior, respectively, which helped shape development of several treatment sessions; Sarah Pearlstein and Sarah Dowling for conducting diagnostic interviews and overseeing project management; Karalani Cross and Taylor Smith for overseeing project management; and Carl Bolano, Kevin Carlis, Michelle Chang, Joanna Chen, Melody Chen, Christina Cui, Vivi Dang, Angelica Estrada, Alyson Johnson, Sanskruti Kakaria, Sarah Knapp, Stephanie Lee, Mercy Lopez, Gregory Pak, Jasmine Rai, Atiyeh Samadi, Rachel Storer, Aaron Tay, Sarah Tran, Stephanie Zepeda for their help with recruitment, screening, data collection and management.

Table 1

Positive Activity Intervention Protocol.

Module	Goal	Description
Psychoeducation Session 1	Present treatment rationale; self-monitoring of emotions.	Present model of emotions (downward spiral of negative emotions; upward spiral of positive emotions). Discuss the nature and function of positive emotions, and how anxiety and depression can disrupt positive experiences.
Noticing & Amplifying Positive Events Session 1, Exercise 1a Session 2, Exercise 1b	Train attention towards and increase awareness of positive events; intensify and prolong positive emotional experiences.	Participants monitor and note positive events throughout the week, and implement strategies to intensify and prolong positive emotions (e.g., savoring, reminiscing, writing about positive events, sharing positive events with others) ^{117, 118} .
Gratitude: Counting One's Blessings Session 2; Exercise 2a Session 3; Exercise 2b	Direct attention to positive events.	Participants write about up to five things for which they are grateful, including events from the past week, or things in their life more generally ^{99, 119, 120} .
Acts of Kindness Session 3; Exercise 3a Session 4; Exercise 3b	Increase prosocial behaviors.	Participants perform up to five kind acts for others within a single day ^{100, 121, 122} .
Pleasurable, engaging, and meaningful activities Session 4; Exercise 4a Session 5; Exercise 4b	Increase participation in rewarding activities, both hedonic and eudaimonic.	Participants complete at least one pleasurable activity alone, one pleasurable activity with others, one highly engaging activity, and one meaningful or important activity ^{123, 124} .
Strengths	Increase awareness of	Participants identify opportunities to use at least one strength

Session 5; Exercise 5a Session 5; Exercise 5b	personal strengths and opportunities to use those strengths.	each day, and monitor the outcome ⁹⁹ .
Affirming Values Session 5; Exercise 5c	Strengthen commitment to personal values.	Participants identify a personal value (i.e., something that makes their life precious and worthwhile) and write about why that value is important to them, and how they use that value in their everyday life ¹²⁵ .
Optimism: Best Possible Future Session 6; Exercise 6a Session 7; Exercise 6b	Promote future-oriented positive cognitions and behaviors.	Participants imagine one's best possible future in chosen domain (e.g., school/career, social relationships, health), and consider how to actualize this future ¹²⁶⁻¹²⁸ .
Make Someone Else Happier Session 7; Exercise 7a Session 8; Exercise 7b	Increase investment in relationships; increase prosocial behaviors and connection with others.	Participants invest time and effort throughout the week into making someone else happier ^{129, 130} .
Live this Month Like It's Your Last in this Area Session 8; Exercise 8a Session 9; Exercise 8b	Promote increased engagement in positive activities; increase anticipation, appreciation, and sustained responsiveness to positive events.	Participants consider what they like most about their current surroundings (e.g., special people, places, activities) and take part in activities as though they will be moving far away within the next month. Participants amplify those experiences through savoring and appreciation ¹³¹ .
Gratitude: Gratitude Letter Session 9; Exercise 9a	Direct attention to another person's role in positive events; strengthen social connections.	Participants recall another person's kindness and write a letter that describes their gratitude (the letter can be sent if desired) ^{99, 126, 128} .
Develop Personalized	Maximize person-activity	Participants develop a positive activity plan by selecting

Positive Activity Plan Session 9; Exercise 9b	fit and continued engagement in activities; translate activities into habits.	activities that are the best fit (e.g., enjoyable, beneficial), and personalizing activities for continued improvement and lasting changes ¹¹⁰ .
Termination Plan Session 10; Exercise 10	Consolidate learning; develop plan to maintain/enhance gains.	Review material and exercises; identify lessons learned and most beneficial exercises; develop plan for continued commitment to engage in these activities.

Table 2

Patient Demographic and Clinical Characteristics

Variable	PAI (n=16)	WL (n=12)
Gender (% female)	50%	75%
Age	29.8 (12.2)	29.0 (12.0)
Years of Education	16.8 (2.7)	15.3 (2.9)
Race (%)		
Caucasian	75%	67%
Asian American	19%	25%
Native American	6%	0%
Pacific Islander	0%	8%
Hispanic (%)	19%	25%
Diagnoses (%)		
Major Depressive Disorder	56%	56%
Social Anxiety Disorder	56%	58%
Generalized Anxiety Disorder	31%	50%
Posttraumatic Stress Disorder	13%	33%
Panic Disorder	0%	17%
Obsessive Compulsive Disorder	0%	8%
Eating Disorder	0%	25%
Mild Alcohol Use Disorder	6%	8%
Mild Marijuana Use Disorder	0%	8%
Past Psychotropic Medication Use (%)	38%	50%
Past Psychosocial Treatment (%)	88%	50%

Note. Standard deviations in parentheses. Percentages sum to > 100% given high comorbidity across the sample.

Table 3

Descriptive summaries of the treatment outcome measures for the PAI (n=16) and waitlist (n=12) groups.

Measure	Pre-assessment <i>M (SD)</i>		Post-assessment <i>M (SD)</i>		Results ^a (Group)		
	PAI	WL	PAI	WL	<i>F</i>	<i>p</i>	η^2_p
Positive Emotions Composite	-0.41 (.96)	-0.40 (.54)	0.73 (.87)	-0.02 (.79)	6.89	.015	.22
PANAS-PA	23.50 (8.43)	24.25 (5.79)	32.63 (7.36)	27.42 (6.87)			
mDES-Positive emotions	15.56 (8.04)	15.08 (3.20)	24.31 (7.73)	17.75 (6.41)			
Negative Emotions Composite	0.17 (.62)	0.70 (.89)	-0.91 (.40)	0.29 (1.11)	10.68	.003	.30
PANAS-NA	25.25 (7.04)	30.17 (9.43)	15.13 (3.56)	26.00 (9.53)			
mDES-Negative emotions	18.81 (6.83)	23.83 (7.85)	8.56 (4.59)	20.25 (12.02)			
Psychological Well-Being Composite	-0.25 (.77)	-0.34 (.81)	0.76 (.81)	-0.35 (.87)	16.91	.000	.40
SWLS	14.75 (6.94)	14.25 (6.41)	22.69 (6.83)	13.67 (6.68)			
QLES-Q	38.06 (7.46)	36.92 (9.22)	47.75 (9.75)	37.50 (9.65)			
Anxiety Symptoms Composite	0.14 (.54)	0.40 (.62)	-0.61 (.68)	0.23 (.91)	5.97	.022	.19
OASIS	10.89 (3.05)	10.33 (4.36)	5.25 (3.00)	8.92 (4.83)			
STAI-Trait	45.00 (7.25)	49.33 (2.67)	43.27 (7.47)	49.25 (6.34)			
Depression Symptoms Composite	0.29 (.90)	0.47 (.77)	-0.87 (.55)	0.29 (.97)	16.01	.000	.39
BDI-II	24.19 (11.73)	27.33 (9.59)	7.73 (5.79)	24.58 (14.38)			
PHQ-9	12.19 (5.84)	12.92 (5.09)	5.81 (4.04)	12.08 (5.25)			

Note. PAI = Positive Activity Intervention; WL = Waitlist; PANAS = Positive and Negative Affect Schedule; PA = positive affect;

NA = negative affect; mDES = Modified Differential Emotions Scale; SWLS = Satisfaction with Life Scale; Q-LES-Q = Quality of

Life Enjoyment and Satisfaction Questionnaire; OASIS = Overall Anxiety and Severity and Impairment Scale; STAI = Spielberger State-Trait Anxiety Inventory (STAI); BDI-II = Beck Depression Inventory II; PHQ-9 = Patient Health Questionnaire.

^a Analysis of covariance (ANCOVA) results comparing the PAI and WL groups at post-assessment controlling for pre-assessment dependent outcomes.

Table 4

Effect sizes for pre- to post-treatment change in the PAI and waitlist groups.

Measure	Effect Size		
	Within-group ^a		Between-groups ^b
	PAI	WL	
Positive Emotions Composite	1.25	0.57	1.00
PANAS-PA	1.16	0.50	0.87
mDES-Positive emotions	1.11	0.56	0.90
Negative Emotions Composite	-2.12	-0.41	-1.28
PANAS-NA	-1.91	-0.44	-1.43
mDES-Negative emotions	-1.80	-0.36	-1.12
Psychological Well-Being Composite	1.28	-0.01	1.57
SWLS	1.15	-0.09	1.73
QLES-Q	1.13	0.06	1.13
Anxiety Composite	-1.23	-0.22	-0.94
OASIS	-1.86	-0.31	-1.01
STAI-Trait	-0.24	-0.02	-0.48
Depression Composite	-1.60	-0.21	-1.54
BDI-II	-1.88	-0.23	-1.53
PHQ-9	-1.29	-0.16	-1.40

Note. PAI = Positive Activity Intervention; WL = Waitlist; PANAS = Positive and Negative Affect Schedule; PA = positive affect; NA = negative affect; mDES = Modified Differential Emotions Scale; SWLS = Satisfaction with Life Scale; Q-LES-Q = Quality of Life Enjoyment and Satisfaction Questionnaire; OASIS = Overall Anxiety and Severity and Impairment Scale; STAI = Spielberger State-Trait Anxiety Inventory (STAI); BDI-II = Beck Depression Inventory II; PHQ-9 = Patient Health Questionnaire.

^aWithin-group pre- to post-treatment effect sizes = $([\text{post-treatment mean} - \text{pre-treatment mean}] / [\text{pre-treatment standard deviation} + \text{post-treatment standard deviation}] / 2)$.

^bBetween-group controlled effect sizes = (post-assessment PAI group covariance adjusted mean – post-assessment waitlist group covariance adjusted mean)/pooled standard deviation.

Table 5

Descriptive summaries of the treatment outcome measures at pre, post, and follow-up assessments for the PAI group (n=15).

Measure	Pre- assessment <i>M (SD)</i>	Post- assessment <i>M (SD)</i>	3-month follow-up <i>M (SD)</i>	6-month follow-up <i>M (SD)</i>	Results ^a (Time)		
					<i>F</i>	<i>p</i>	η^2_p
Positive Emotions Composite	-0.79 (.91)	0.26 (.82)	0.31 (.73)	0.21 (.85)	14.58	.000	.51
PANAS-PA	23.50 (8.43)	32.63 (7.36)	35.19 (6.22)	33.06 (7.11)			
mDES-Positive emotions	15.56 (8.04)	24.31 (7.73)	22.94 (7.40)	23.50 (8.07)			
Negative Emotions Composite	1.05 (.86)	-0.54 (.61)	-0.28 (.50)	-0.23 (.80)	19.10	.000	.58
PANAS-NA	25.25 (7.04)	15.13 (3.56)	17.19 (2.99)	17.50 (5.67)			
mDES-Negative emotions	18.81 (6.83)	8.56 (4.59)	9.19 (3.49)	9.44 (4.19)			
Psychological Well-Being Composite	-0.77 (.79)	0.22 (.84)	0.26 (.75)	0.28 (.86)	23.28	.000	.62
SWLS	14.75 (6.94)	22.69 (6.83)	22.94 (7.62)	23.06 (7.30)			
QLES-Q	38.06 (7.46)	47.75 (9.75)	48.06 (6.21)	48.25 (9.55)			
Anxiety Symptoms Composite	0.51 (.56)	-0.27 (.69)	-0.14 (.66)	-0.11 (.57)	9.21	.009	.40
OASIS	10.89 (3.05)	5.25 (3.00)	5.19 (4.10)	5.25 (4.20)			
STAI-Trait	45.00 (7.25)	43.27 (7.47)	44.75 (5.50)	45.00 (5.02)			
Depression Symptoms Composite	0.99 (1.04)	-0.33 (.64)	-0.32 (.59)	-0.33 (.78)	18.29	.000	.57
BDI-II	24.19 (11.73)	7.73 (5.79)	9.00 (6.63)	8.44 (8.79)			
PHQ-9	12.19 (5.84)	5.81 (4.04)	4.88 (3.52)	5.06 (4.11)			

Note. PAI = Positive Activity Intervention; WL = Waitlist; PANAS = Positive and Negative Affect Schedule; PA = positive affect;

NA = negative affect; mDES = Modified Differential Emotions Scale; SWLS = Satisfaction with Life Scale; Q-LES-Q = Quality of

Life Enjoyment and Satisfaction Questionnaire; OASIS = Overall Anxiety and Severity and Impairment Scale; STAI = Spielberger State-Trait Anxiety Inventory (STAI); BDI-II = Beck Depression Inventory II; PHQ-9 = Patient Health Questionnaire.

^a Repeated measures analysis of variance (ANOVA) results for Time (pre-, post-, 3-month, 6-month follow-up) in the PAI group.

Figure Legends

Figure 1. CONSORT flow diagram illustrating participants' progress throughout the study.

References

1. Kessler RC, Petukhova M, Sampson NA, Zaslavsky AM, Wittchen HU. Twelve-month and lifetime prevalence and lifetime morbid risk of anxiety and mood disorders in the United States. *Int J Methods Psychiatr Res* 2012;21(3):169-84.
2. Baxter AJ, Vos T, Scott KM, Ferrari AJ, Whiteford HA. The global burden of anxiety disorders in 2010. *Psychol Med* 2014;44(11):2363-74.
3. Whiteford HA, Degenhardt L, Rehm J, Baxter AJ, Ferrari AJ, Erskine HE, et al. Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. *Lancet* 2013;382(9904):1575-86.
4. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005;62(6):617-27.
5. Rapaport MH, Clary C, Fayyad R, Endicott J. Quality-of-life impairment in depressive and anxiety disorders. *Am J Psychiatry* 2005;162(6):1171-8.
6. Olatunji BO, Cisler JM, Tolin DF. Quality of life in the anxiety disorders: a meta-analytic review. *Clin Psychol Rev* 2007;27(5):572-81.
7. Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, et al. Research domain criteria (RDoC): Toward a new classification framework for research on mental disorders. *Am J Psychiatry* 2010;167(7):748-51.
8. Dillon DG, Rosso IM, Pechtel P, Killgore WDS, Rauch SL, Pizzagalli DA. Peril and pleasure: An RDoC-inspired examination of threat responses and reward processing in anxiety and depression. *Depress Anxiety* 2014;31(3):233-49.

9. Fredrickson BL. Positive emotions broaden and build. *Adv Exp Soc Psychol* 2013;47:1-53.
10. Lyubomirsky S, King L, Diener E. The benefits of frequent positive affect: Does happiness lead to success? *Psychol Bull* 2005;131(6):803-55.
11. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (5th Ed.). *Author* Washington, DC 2013.
12. Barlow DH, Sauer-Zavala S, Carl JR, Bullis JR, Ellard KK. The nature, diagnosis, and treatment of neuroticism: Back to the future. *Clin Psychol Sci* 2014;2(3):344-65.
13. Barlow DH, Allen LB, Choate ML. Toward a unified treatment for emotional disorders. *Behav Ther* 2004;35(2):205-30.
14. Hofmann SG, Smits JA. Cognitive-behavioral therapy for adult anxiety disorders: a meta-analysis of randomized placebo-controlled trials. *J Clin Psychiatry* 2008;69(4):621-32.
15. Norton PJ, Price EC. A meta-analytic review of adult cognitive-behavioral treatment outcome across the anxiety disorders. *J Nerv Ment Dis* 2007;195(6):521-31.
16. Craske MG, Kircanski K, Zelikowsky M, Mystkowski J, Chowdhury N, Baker A. Optimizing inhibitory learning during exposure therapy. *Behav Res Ther* 2008;46(1):5-27.
17. Craske MG, Treanor M, Conway CC, Zbozinek T, Vervliet B. Maximizing exposure therapy: an inhibitory learning approach. *Behav Res Ther* 2014;58:10-23.
18. Cuijpers P, Berking M, Andersson G, Quigley L, Kleiboer A, Dobson KS. A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments. *Can J Psychiatry* 2013;58(7):376-85.
19. Dunn BD. Helping depressed clients reconnect to positive emotion experience: Current insights and future directions. *Clin Psychol Psychother* 2012;19(4):326-40.

20. Garland EL, Fredrickson B, Kring AM, Johnson DP, Meyer PS, Penn DL. Upward spirals of positive emotions counter downward spirals of negativity: Insights from the broaden-and-build theory and affective neuroscience on the treatment of emotion dysfunctions and deficits in psychopathology. *Clin Psychol Rev* 2010;30(7):849-64.
21. Layous K, Chancellor J, Lyubomirsky S. Positive activities as protective factors against mental health conditions. *J Abnorm Psychol* 2014;123(1):3-12.
22. Fredrickson BL, Levenson RW. Positive emotions speed recovery from the cardiovascular sequelae of negative emotions. *Cogn Emot* 1998;12(2):191-220.
23. Fredrickson BL, Mancuso RA, Branigan C, Tugade MM. The undoing effect of positive emotions. *Motiv Emot* 2000;24(4):237-58.
24. Fredrickson BL, Tugade MM, Waugh CE, Larkin GR. What good are positive emotions in crises? A prospective study of resilience and emotions following the terrorist attacks on the United States on September 11th, 2001. *J Pers Soc Psychol* 2003;84(2):365-76.
25. Tugade MM, Fredrickson BL. Resilient individuals use positive emotions to bounce back from negative emotional experiences. *J Pers Soc Psychol* 2004;86(2):320-33.
26. Cunningham MR. What do you do when you're happy or blue: Mood, expectancies, and behavioral interest. *Motiv Emot* 1988;12(4):309-31.
27. Fredrickson BL, Branigan C. Positive emotions broaden the scope of attention and thought-action repertoires. *Cogn Emot* 2005;19(3):313-32.
28. Wadlinger HA, Isaacowitz DM. Positive mood broadens visual attention to positive stimuli. *Motiv Emot* 2006;30(1):89-101.
29. Isen AM. An influence of positive affect on decision making in complex situations: Theoretical issues with practical implications. *J Consum Psychol* 2001;11(2):75-85.

30. Lyubomirsky S, Boehm JK, Kasri F, Zehm K. The cognitive and hedonic costs of dwelling on achievement-related negative experiences: Implications for enduring happiness and unhappiness. *Emotion* 2011;11(5):1152-67.
31. Gable SL, Reis HT, Elliot AJ. Behavioral activation and inhibition in everyday life. *J Pers Soc Psychol* 2000;78(6):1135-49.
32. Gable SL, Berkman ET. Making connections and avoiding loneliness: Approach and avoidance social motives and goals. In: Elliot AJ, editor. Handbook of approach and avoidance motivation. New York, NY: Psychology Press; 2008. p. 203-16.
33. Fredrickson BL, Cohn MA, Coffey KA, Pek J, Finkel SM. Open hearts build lives: Positive emotions, induced through Loving-Kindness Meditation, build consequential personal resources. *J Pers Soc Psychol* 2008;95(5):1045-62.
34. Keyes CL. Mental illness and/or mental health? Investigating axioms of the complete state model of health. *J Consult Clin Psychol* 2005;73(3):539.
35. Kuppens P, Realo A, Diener E. The role of positive and negative emotions in life satisfaction judgment across nations. *J Pers Soc Psychol* 2008;95(1):66-75.
36. Kashdan TB. Social anxiety spectrum and diminished positive experiences: theoretical synthesis and meta-analysis. *Clin Psychol Rev* 2007;27(3):348-65. Epub 2007/01/16.
37. Kashdan TB, Weeks JW, Savostyanova AA. Whether, how, and when social anxiety shapes positive experiences and events: A self-regulatory framework and treatment implications. *Clin Psychol Rev* 2011;31(5):786-99.
38. Nawijn L, van Zuiden M, Frijling JL, Koch SBJ, Veltman DJ, Olff M. Reward functioning in PTSD: A systematic review exploring the mechanisms underlying anhedonia. *Neurosci Biobehav Rev* 2015;51:189-204.

39. Brown TA, Chorpita BF, Barlow DH. Structural relationships among dimensions of the DSM-IV anxiety and mood disorders and dimensions of negative affect, positive affect, and autonomic arousal. *J Abnorm Psychol* 1998;107(2):179-92.
40. Naragon-Gainey K, Watson D, Markon KE. Differential relations of depression and social anxiety symptoms to the facets of extraversion/positive emotionality. *J Abnorm Psychol* 2009;118(2):299-310.
41. Watson D, Naragon-Gainey K. On the specificity of positive emotional dysfunction in psychopathology: evidence from the mood and anxiety disorders and schizophrenia/schizotypy. *Clin Psychol Rev* 2010;30(7):839-48.
42. Campbell DW, Sareen J, Stein MB, Kravetsky LB, Paulus MP, Hassard ST, et al. Happy but not so approachable: the social judgments of individuals with generalized social phobia. *Depress Anxiety* 2009;26(5):419-24.
43. Clausen AN, Youngren W, Sisante JF, Billinger SA, Taylor C, Aupperle RL. Combat PTSD and implicit behavioral tendencies for positive affective stimuli: A brief report. *Front Psychol* 2016;7:758.
44. Sherdell L, Waugh CE, Gotlib IH. Anticipatory pleasure predicts motivation for reward in major depression. *J Abnorm Psychol* 2012;121(1):51-60.
45. Heuer K, Rinck M, Becker ES. Avoidance of emotional facial expressions in social anxiety: The Approach-Avoidance Task. *Behav Res Ther* 2007;45(12):2990-3001.
46. Trew JL. Exploring the roles of approach and avoidance in depression: an integrative model. *Clin Psychol Rev* 2011;31(7):1156-68.
47. Joormann J, Gotlib IH. Selective attention to emotional faces following recovery from depression. *J Abnorm Psychol* 2007;116(1):80-5.

48. Joormann J, Talbot L, Gotlib IH. Biased processing of emotional information in girls at risk for depression. *J Abnorm Psychol* 2007;116(1):135-43.
49. Taylor CT, Bomyea J, Amir N. Malleability of attentional bias for positive emotional information and anxiety vulnerability. *Emotion* 2011;11(1):127-38.
50. Taylor CT, Bomyea J, Amir N. Attentional bias away from positive social information mediates the link between social anxiety and anxiety vulnerability to a social stressor. *J Anxiety Disord* 2010;24(4):403-8.
51. Pizzagalli DA, Iosifescu D, Hallett LA, Ratner KG, Fava M. Reduced hedonic capacity in major depressive disorder: Evidence from a probabilistic reward task. *J Psychiatr Res* 2008;43(1):76-87.
52. Pizzagalli DA, Holmes AJ, Dillon DG, Goetz EL, Birk JL, Bogdan R, et al. Reduced caudate and nucleus accumbens response to rewards in unmedicated individuals with major depressive disorder. *Am J Psychiatry* 2009;166(6):702-10.
53. Sailer U, Robinson S, Fischmeister FPS, Konig D, Oppenauer C, Lueger-Schuster B, et al. Altered reward processing in the nucleus accumbens and mesial prefrontal cortex of patients with posttraumatic stress disorder. *Neuropsychologia* 2008;46(11):2836-44.
54. Elman I, Lowen S, Frederick BB, Chi W, Becerra L, Pitman RK. Functional Neuroimaging of reward circuitry responsivity to monetary gains and losses in posttraumatic stress disorder. *Biol Psychiatry* 2009;66(12):1083-90.
55. Richey JA, Rittenberg A, Hughes L, Damiano CR, Sabatino A, Miller S, et al. Common and distinct neural features of social and non-social reward processing in autism and social anxiety disorder. *Soc Cogn Affect Neurosci* 2014;9(3):367-77.

56. Carl JR, Soskin DP, Kerns C, Barlow DH. Positive emotion regulation in emotional disorders: A theoretical review. *Clin Psychol Rev* 2013;33(3):343-60.
57. Davidson RJ. Affective style and affective disorders: Perspectives from affective neuroscience. *Cogn Emot* 1998;12(3):307-30.
58. Davidson RJ, Jackson DC, Kalin NH. Emotion, plasticity, context, and regulation: Perspectives from affective neuroscience. *Psychol Bull* 2000;126(6):890-909.
59. Mazzucchelli TG, Kane RT, Rees CS. Behavioral activation interventions for well-being: A meta-analysis. *J Posit Psychol* 2010;5(2):105-21.
60. Forman EM, Herbert JD, Moitra E, Yeomans PD, Geller PA. A randomized controlled effectiveness trial of acceptance and commitment therapy and cognitive therapy for anxiety and depression. *Behav Modif* 2007;31(6):772-99.
61. Oei TP, McAlinden NM. Changes in quality of life following group CBT for anxiety and depression in a psychiatric outpatient clinic. *Psychiatry Res* 2014;220(3):1012-8.
62. Safren SA, Heimberg RG, Brown EJ, Holle C. Quality of life in social phobia. *Depress Anxiety* 1996;4(3):126-33.
63. Kring AM, Persons JB, Thomas C. Changes in affect during treatment for depression and anxiety. *Behav Res Ther* 2007;45(8):1753-64.
64. Dunn BD, German RE, Hollon SD, DeRubeis RJ. Changes in positive and negative affect during cognitive therapy versus medication for major depressive disorder: A randomized controlled trial. *Manuscript submitted for publication* 2016.
65. Lyubomirsky S, Sheldon KM, Schkade D. Pursuing happiness: The architecture of sustainable change. *Rev Gen Psychol* 2005;9(2):111-31.

66. Sin NL, Lyubomirsky S. Enhancing well-being and alleviating depressive symptoms with positive psychology interventions: A practice-friendly meta-analysis. *J Clin Psychol* 2009;65(5):467-87.
67. Bolier L, Haverman M, Westerhof GJ, Riper H, Smit F, Bohlmeijer E. Positive psychology interventions: a meta-analysis of randomized controlled studies. *BMC Public Health* 2013;13.
68. Seligman MEP, Rashid T, Parks AC. Positive psychotherapy. *Am Psychol* 2006;61(8):774-88.
69. Huffman JC, Mastromauro CA, Boehm JK, Seabrook R, Fricchione GL, Denninger JW, et al. Development of a positive psychology intervention for patients with acute cardiovascular disease. *Heart Int* 2011;6(2):e14.
70. Huffman JC, DuBois CM, Healy BC, Boehm JK, Kashdan TB, Celano CM, et al. Feasibility and utility of positive psychology exercises for suicidal inpatients. *Gen Hosp Psychiat* 2014;36(1):88-94.
71. Moskowitz JT, Hult JR, Duncan LG, Cohn MA, Maurer S, Bussolari C, et al. A positive affect intervention for people experiencing health-related stress: development and non-randomized pilot test. *J Health Psychol* 2012;17(5):676-92.
72. Campbell-Sills L, Norman SB, Craske MG, Sullivan G, Lang AJ, Chavira DA, et al. Validation of a brief measure of anxiety-related severity and impairment: the Overall Anxiety Severity and Impairment Scale (OASIS). *J Affect Disord* 2009;112(1-3):92-101.
73. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9 - Validity of a brief depression severity measure. *J Gen Intern Med* 2001;16(9):606-13.

74. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: The PANAS scales. *J Pers Soc Psychol* 1988;54(6):1063-70.
75. Endicott J, Nee J, Harrison W, Blumenthal R. Quality of Life Enjoyment and Satisfaction Questionnaire: A new measure. *Psychopharmacol Bull* 1993;29(2):321-6.
76. Stevanovic D. Quality of Life Enjoyment and Satisfaction Questionnaire-short form for quality of life assessments in clinical practice: A psychometric study. *J Psychiatr Ment Health Nurs* 2011;18(8):744-50.
77. Diener E, Emmons RA, Larsen RJ, Griffin S. The Satisfaction With Life Scale. *J Pers Assess* 1985;49(1):71-5.
78. Pavot W, Diener E, Colvin CR, Sandvik E. Further validation of the Satisfaction with Life Scale: Evidence for the cross-method convergence of well-being measures. *J Pers Assess* 1991;57(1):149-61.
79. Pavot W, Diener E. Review of the Satisfaction with Life Scale. In: E. Diener, Editor. *Assessing well-being*. Netherlands: Springer; 2009. p. 101-17.
80. Norman SB, Cissell SH, Means-Christensen AJ, Stein MB. Development and validation of an Overall Anxiety Severity and Impairment Scale (OASIS). *Depress Anxiety* 2006;23(4):245-9.
81. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA. *Manual for the State-Trait Anxiety Inventory*. Palo Alto, CA: Consulting Psychologist Press; 1983.
82. Kroenke K, Spitzer RL, Williams JB. A new measure of depression severity: The PHQ-9. *J Gen Intern Med* 2000;15:78.
83. Beck AT, Steer, A., & Brown K. *Beck Depression Inventory-II*. 1996.

84. Dozois DJA, Dobson, K. S., & Ahnberg, J. L. A psychometric evaluation of the Beck Depression Inventory - II. *Psychol Assess* 1998; 10:83-89.
85. Devilly GJ, Borkovec TD. Psychometric properties of the credibility/expectancy questionnaire. *J Behav Ther Exp Psy* 2000;31(2):73-86.
86. Taylor CT, Cissell SH, Lyubomirsky S. Positive activity intervention for anxiety and depression: Treatment manual. 2014.
87. Lyubomirsky S, Layous K. How do simple positive activities increase well-being? *Curr Dir Psychol Sci* 2013;22(1):57-62.
88. Layous K, Chancellor J, Lyubomirsky S, Wang L, Doraiswamy PM. Delivering happiness: Translating positive psychology intervention research for treating major and minor depressive disorders. *J Altern Complement Med* 2011;17(8):675-83.
89. Fredrickson BL. What good are positive emotions? *Rev Gen Psychol* 1998;2(3):300-19.
90. Fredrickson BL. The value of positive emotions - The emerging science of positive psychology in coming to understand why it's good to feel good. *Am Sci* 2003;91(4):330-5.
91. Fredrickson BL. The role of positive emotions in positive psychology - The broaden-and-build theory of positive emotions. *Am Psychol* 2001;56(3):218-26.
92. Van Breukelen GJP. ANCOVA versus change from baseline had more power in randomized studies and more bias in nonrandomized studies. *J Clin Epidemiol* 2006;59:920-5.
93. Vickers AJ, Altman DG. Statistics notes: Analysing controlled trials with baseline and follow up measurements. *BMJ* 2001;323(7321):1123-4.
94. Clark DM, Ehlers A, Hackmann A, McManus F, Fennell M, Grey N, et al. Cognitive therapy versus exposure and applied relaxation in social phobia: A randomized controlled trial. *J Consult Clin Psychol* 2006;74(3):568-78.

95. Clark DM, Ehlers A, McManus F, Hackmann A, Fennell M, Campbell H, et al. Cognitive therapy versus fluoxetine in generalized social phobia: a randomized placebo-controlled trial. *J Consult Clin Psychol* 2003;71(6):1058-67.
96. Hollon SD, DeRubeis RJ, Evans MD, Wiemer MJ, Garvey MJ, Grove WM, et al. Cognitive therapy and pharmacotherapy for depression. Singly and in combination. *Arch Gen Psychiatry* 1992;49(10):774-81.
97. Rosenthal R, Rosnow RL. Essentials of behavioral research: Methods and data analysis (2nd ed.). New York: McGraw-Hill; 1991.
98. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Front Psychol* 2013;4:863.
99. Seligman MEP, Steen TA, Park N, Peterson C. Positive psychology progress - Empirical validation of interventions. *Am Psychol* 2005;60(5):410-21.
100. Alden LE, Trew JL. If It Makes You Happy: Engaging in kind acts increases positive affect in socially anxious individuals. *Emotion* 2013;13(1):64-75.
101. Crawford JR, Henry JD. The positive and negative affect schedule (PANAS): Construct validity, measurement properties and normative data in a large non-clinical sample. *Brit J Clin Psychol* 2004;43:245-65.
102. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: The PANAS scales. *J Pers Soc Psychol* 1988;54(6):1063-70.
103. Pavot W, Diener E. The affective and cognitive context of self-reported measures of subjective well-being. *Soc Indic Res* 1993;28(1):1-20.

104. Norton PJ, Barrera TL. Transdiagnostic versus diagnosis-specific cbt for anxiety disorders: a preliminary randomized controlled noninferiority trial. *Depress Anxiety* 2012;29(10):874-82.
105. Farchione TJ, Fairholme CP, Ellard KK, Boisseau CL, Thompson-Hollands J, Carl JR, et al. Unified protocol for transdiagnostic treatment of emotional disorders: A randomized controlled trial. *Behav Ther* 2012;43(3):666-78.
106. Mansell W, Harvey A, Watkins E, Shafran R. Conceptual foundations of the transdiagnostic approach to CBT. *J Cogn Psychother* 2009;23(1):6-19.
107. Jacobson NS, Truax P. Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *J Consult Clin Psychol* 1991;59(1):12-19.
108. Maas CJM, Hox JJ. Sufficient sample sizes for multilevel modeling. *Methodology* 2005;1(3):86-92.
109. Devilly GJ, McFarlane AC. When wait lists are not feasible, nothing is a thing that does not need to be done. *J Consult Clin Psychol* 2009;77(6):1159-1168.
110. Nelson SK, Lyubomirsky S. Finding happiness: Tailoring positive activities for optimal well-being benefits. In: Tugade M, Shiota M, Kirby L, Editors. Handbook of positive emotions. New York: Guilford; 2014. p. 275-93.
111. Ozomaro U, Wahlestedt C, Nemeroff CB. Personalized medicine in psychiatry: problems and promises. *BMC Med* 2013;11.
112. Gable SL, Reis HT, Elliot AJ. Evidence for bivariate systems: An empirical test of appetite and aversion across domains. *J Res Pers* 2003;37(5):349-72.
113. Lang PJ. The emotion probe. Studies of motivation and attention. *Am Psychol* 1995;50(5):372-85.

114. Gray JA. The psychology of fear and stress (2nd ed.). New York: Cambridge University Press; 1987.
115. Elliot AJ. The hierarchical model of approach-avoidance motivation. *Motiv Emot* 2006;30(2):111-6.
116. Elliot AJ, Thrash TM. Approach and avoidance temperament as basic dimensions of personality. *J Pers* 2010;78(3):865-906.
117. Jose PE, Lim BT, Bryant FB. Does savoring increase happiness? A daily diary study. *J Posit Psychol* 2012;7(3):176-87.
118. McMakin DL, Siegle GJ, Shirk SR. Positive affect stimulation and sustainment (PASS) module for depressed mood: A preliminary investigation of treatment-related effects. *Cognit Ther Res* 2011;35(3):217-26.
119. Chancellor J, Layous K, Lyubomirsky S. Recalling positive events at work makes employees feel happier, move more, but interact less: A 6-week randomized controlled intervention at a Japanese workplace. *J Happiness Stud* 2015;16(4):871-87.
120. Emmons RA, McCullough ME. Counting blessings versus burdens: An experimental investigation of gratitude and subjective well-being in daily life. *J Pers Soc Psychol* 2003;84(2):377-89.
121. Layous K, Nelson SK, Oberle E, Schonert-Reichl KA, Lyubomirsky S. Kindness counts: Prompting prosocial behavior in preadolescents boosts peer acceptance and well-being. *PLoS One* 2012;7(12).
122. Nelson SK, Layous K, Cole S, Lyubomirsky S. Do unto others or treat yourself? The effects of prosocial and self-focused behavior on psychological flourishing. *Emotion* in press.

123. Park N, Peterson C, Ruch W. Orientations to happiness and life satisfaction in twenty-seven nations. *J Posit Psychol* 2009;4(4):273-9.
124. Csikszentmihalyi M. The contribution of flow to positive psychology. *Laws Lif Symp Ser* 2000;2:387-95.
125. Nelson SK, Fuller JAK, Choi I, Lyubomirsky S. Beyond self-protection: Self-affirmation benefits hedonic and eudaimonic well-being. *Pers Soc Psychol B* 2014;40(8):998-1011.
126. Boehm JK, Lyubomirsky S, Sheldon KM. A longitudinal experimental study comparing the effectiveness of happiness-enhancing strategies in Anglo Americans and Asian Americans. *Cogn Emot* 2011;25(7):1263-72.
127. Layous K, Nelson SK, Lyubomirsky S. What is the optimal way to deliver a positive activity intervention? The case of writing about one's best possible selves. *J Happiness Stud* 2013;14(2):635-54.
128. Lyubomirsky S, Dickerhoof R, Boehm JK, Sheldon KM. Becoming happier takes both a will and a proper way: An experimental longitudinal intervention to boost well-being. *Emotion* 2011;11(2):391-402.
129. Dunn EW, Aknin LB, Norton MI. Spending money on others promotes happiness. *Science* 2008;319(5870):1687-8.
130. Layous K, Kurtz J, Chancellor J, Lyubomirsky S. Make someone happy ... and you will be happy too: An other-oriented path to well-being. *Manuscript submitted for publication*. 2016.
131. Layous K, Kurtz J, Chancellor J, Lyubomirsky S. Reframing the ordinary: Imagining time as scare increases well-being. *Manuscript submitted for publication*. 2016.